arXiv:1403.6101v1 [math.AT] 24 Mar 2014

Homotopy theory of G-diagrams and equivariant excision

Emanuele Dotto and Kristian Moi

March 25, 2014

Abstract

Let G be a finite group acting on a small category I. We study functors $X\colon I \to \mathscr{C}$ equipped with families of compatible natural transformations that give a kind of generalized G-action on X. Such objects are called G-diagrams. When $\mathscr{C}$ is a sufficiently nice model category we give a model structure on the category of I-indexed G-diagrams in $\mathscr{C}$. There are natural G-actions on Bousfield-Kan style homotopy limits and colimits of G-diagrams. We prove that weak equivalences between point-wise (co)fibrant objects in our model structure induce weak G-equivalences on homotopy (co)limits. A case of particular interest is when the indexing category is a cubical diagram. We use homotopy limits and colimits over such diagrams to produce loop and suspension spaces with respect to permutation representations of G. We go on to develop a theory of enriched homotopy functors and give an equivariant linearity condition in terms of cubical G-diagrams, which in the case of G-spaces we prove to be equivalent to Blumberg's notion of G-linearity. In particular we show that the Wirthmüller isomorphism theorem is a direct consequence of the equivariant linearity of the identity functor on G-spectra.

Contents

1 Definitions and setup **4**
1.1 Categories of G-diagrams 4
1.2 Enrichments and homotopy (co)limits 6
1.3 Examples of G-diagrams 10

2 G-diagrams and model structures **11**
2.1 Equivariant model categories 11
2.2 A model structure on G-diagrams 13
2.3 Cofibrant replacement of G-diagrams 16
2.4 Homotopy invariance of map and tensor 19
2.5 Equivariant cofinality 20
2.6 The Elmendorf theorem for G-diagrams 21

3 Equivariant excision **22**
3.1 Equivariant cubes and G-excision 22
3.2 The generalized Wirthmüller isomorphism 27
3.3 G-linearity and adjoint assembly maps 29
3.4 G-linear functors on G-spaces 32
3.5 G-linear functors to G-spectra 33

A Appendix **35**
A.1 Finite categories and cofibrant G-diagrams 35
A.2 Sequential homotopy colimits and finite G-homotopy limits 37

The first author was partially supported by the ERC Adv.Grant No.228082.
The second author was partially supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92)

Introduction

We begin with a motivating example.

Example. Let X be a pointed space with an action of C_2, the cyclic group of order two, with $\sigma\colon X \to X$ representing the action of the non-trivial group element. Assume we have a diagram

$$Y \xrightarrow{f} X \xleftarrow{g} Z \tag{1}$$

of pointed spaces together with mutually inverse homeomorphisms $r\colon Y \to Z$ and $l\colon Z \to Y$ which cover σ in the sense that $f \circ l = \sigma \circ g$ and $g \circ r = \sigma \circ f$. Then the pullback $Y \times_X Z$ inherits a natural C_2-action given by $(y, z) \mapsto (l(z), r(y))$ and similarly, the homotopy pullback

$$Y \times^h_X Z = \{(y, \gamma, z) \in Y \times X^I \times Z \mid f(y) = \gamma(0) \text{ and } \gamma(1) = g(z)\}$$

inherits the action $(y, \gamma, z) \mapsto (l(z), \sigma \circ \bar{\gamma}, r(y))$, where $\bar{\gamma}(t) = \gamma(1 - t)$. The usual inclusion $Y \times_X Z \hookrightarrow Y \times^h_X Z$ is equivariant with respect to these actions. Let $\mathbb{R}^{1,1}$ denote the sign representation of C_2 on $\mathbb{R}$ and let $\Omega^{1,1}X$ be the space of pointed maps from the one point compactification $S^{\mathbb{R}^{1,1}}$ to X with C_2 acting by conjugation. If Y (and hence Z) is contractible, there is a C_2-homotopy equivalence

$$Y \times^h_X Z \simeq \Omega^{1,1}X.$$

On underlying spaces this just an instance of the well-known homotopy equivalence

$$\Omega X \simeq \operatorname{holim}(* \to X \leftarrow *).$$

It is natural to ask when a map between diagrams, such as (1), induces a C_2-homotopy equivalence on homotopy limits and whether one can generalize this to construct loop spaces from other representation spheres.

A structured diagram as in (1) is an instance of a G-diagram. These objects were first introduced independently in [JS01] and [VF99], and were further studied in [VF04] and [TW91] in connection with homotopy colimits of topological spaces and simplicial sets. In the present paper we develop a model categorical framework for G-diagrams, allowing us to deal also with homotopy limit functors, and to work in other categories, for example in the category G-spectra.

Given a finite group G acting on a category I by functors $a(g)\colon I \to I$, a G-diagram in a category $\mathscr{C}$ is a functor $X\colon I \to \mathscr{C}$ together with natural transformations $g_X\colon X \to X \circ a(g)$ for every g in G compatible with the group structure. A map between G-diagram is a natural transformation between the underlying diagrams that commutes with the structure maps (see Definitions 1.1 and 1.2). We write $\mathscr{C}^I_a$ for the category of G-diagrams with these maps as morphisms. The category $\mathscr{C}^I_a$ is isomorphic to the category of diagrams in $\mathscr{C}$ indexed on the Grothendieck construction of the action $a\colon G \to Cat$ (see Lemma 1.8 and [JS01, 2]). If the category of G-objects $\mathscr{C}^G$ is a sufficiently nice model category (for example G-spaces with the fixed points model structure, or orthogonal G-spectra with the G-stable model structure) we prove the following 2.6.

Theorem: *Let $\mathscr{C}$ be a G-model category (see 2.1). There is a cofibrantly generated $sSet^G$-enriched model structure on the category of G-diagrams $\mathscr{C}^I_a$ with*

1. *weak equivalences the maps of G-diagrams $f\colon X \to Y$ whose restrictions at every vertex f_i are equivalences in the model category $\mathscr{C}^{G_i}$ of objects with an action of the stabilizer group G_i of the object i,*
2. *fibrations the maps of G-diagrams $f\colon X \to Y$ whose restrictions at every vertex f_i are fibrations in $\mathscr{C}^{G_i}$ for every object i of I,*

3. generating cofibrations and acyclic cofibrations the collections

$$F\mathcal{I} = \bigcup_{i\in I} F_i\mathcal{I}_i \qquad \text{and} \qquad F\mathcal{J} = \bigcup_{i\in I} F_i\mathcal{J}_i$$

where $\mathcal{I}_i$ and $\mathcal{J}_i$ are respectively generating cofibrations and acyclic cofibrations of $\mathscr{C}^{G_i}$, and $F_i\colon \mathscr{C}^{G_i} \to \mathscr{C}^I_a$ is the left adjoint to the evaluation functor at the vertex i.

When it makes sense to talk about the colimit, limit, homotopy colimit and homotopy limit of a G-diagram X, these constructions have natural G-actions induced by the the structure maps g_X (see Corollary 1.4 and §1.2). Moreover, the usual comparison maps $\lim X \to \operatorname{holim} X$ and $\operatorname{hocolim} X \to \operatorname{colim} X$ are equivariant as we already observed for the C_2-diagram (1). In general most constructions involving (co)limits and (co)ends enrichments applied to G-diagrams produce G-objects and equivariant maps between them. The G-equivariant homotopy type of the Bousfield-Kan homotopy limit and colimit of a G-diagram is homotopy invariant (see 2.22):

Corollary: *The functors* $\operatorname{holim}\colon \mathscr{C}^I_a \to \mathscr{C}^G$ *and* $\operatorname{hocolim}\colon \mathscr{C}^I_a \to \mathscr{C}^G$ *preserve equivalences between fibrant diagrams and pointwise cofibrant diagrams respectively.*

We prove other equivariant analogues of classical theorem of model categories of diagrams, such as a homotopy cofinality theorem for G-diagrams (2.24) generalizing the results [TW91, 1] and [VF04, 6], as well as an Elmendorf-like theorem for G-diagrams (2.25).

As an application of this model categorical theory of G-diagrams, we define and study equivariant excision. Classically, a homotopy invariant functor between model categories is excisive if it sends homotopy cocartesian squares to homotopy cartesian squares. This definition is not well behaved when the categories involved are categories of G-objects. For example, excisive reduced functors from pointed G-spaces to G-spaces are not a model for G-spectra (see [Blu06]) unless G is the trivial group. We formulate G-excision by replacing squares by "equivariant cubes". For a finite G-set J we consider the power set $\mathcal{P}(J)$ as a category with inclusions as morphisms. This has a G-action induced by the G-action on J. A G-diagram indexed on $\mathcal{P}(J)$ is called a J-cube. A J-cube X is defined to be homotopy cocartesian if the map from the homotopy colimit to the value at the terminal vertex is a weak equivalence in $\mathscr{C}^G$. Similarly, it is homotopy cartesian if the map from the value at the initial vertex to the homotopy limit is a weak equivalence in $\mathscr{C}^G$. A suitable homotopy invariant functor $\Phi\colon \mathscr{C}^G \to \mathscr{D}^G$ is called G-excisive if it sends homotopy cocartesian G_+-cubes to homotopy cartesian G_+-cubes. Here G_+ is the set G with a disjoint base point, and G acts on it by left multiplication. It plays the role of a regular G-set, in the same way as the reduced representation of $\mathbb{R}[G_+]$ is the regular representation of G. The paper contains a series of fundamental properties of G-excision, summarized in the following list.

- *3.11 A G-excisive functor $\mathscr{C}^G \to \mathscr{D}^G$ is classically excisive, that is, it sends homotopy cocartesian squares in $\mathscr{C}^G$ to homotopy cartesian squares in $\mathscr{D}^G$,*
- *3.23 For functors on pointed G-spaces $Top^G_* \to Top^G_*$ our definition of G-excision is equivalent to the definition of [Blu06],*
- *3.28 Enriched G-excisive reduced homotopy functors from finite G-CW-complexes to G-spectra are given by smashing with a G-spectrum,*
- *3.27 The identity functor on G-spectra is G-excisive,*
- *3.15 Any G-excisive reduced homotopy functor $\Phi\colon \mathscr{C}^G \to \mathscr{D}^G$ satisfies the Wirthmüller isomorphism theorem, that is, the canonical map $\Phi(G \otimes_H c) \to \hom_H(G, \Phi(c))$ is an equivalence in $\mathscr{D}^G$ for every subgroup H of G and H-object c of $\mathscr{C}$.*
- *3.20,3.21 If $\mathscr{D}^G$ is suitably presentable, a construction similar to Goodwillie's derivative of [Goo92] defines a universal G-excisive approximation to any homotopy functor $\mathscr{C}^G \to \mathscr{D}^G$.*

In particular, the classical Wirthmüller isomorphism theorem for G-spectra follows directly from G-excision for the identity functor on G-spectra by considering a specific homotopy cocartesian cube.

Moreover G-excision for the identity on G-spectra, i.e. the fact that any homotopy cocartesian equivariant J-cube is homotopy cartesian, is a formal consequence of the equivariant Freudenthal suspension theorem. This notion of equivariant cubes and G-excision provides a good context in which the category of G-spectra "is G-stable".

The paper is structured as follows: In section 1 we first define G-diagrams and describe some properties of categories of G-diagrams as well as the interaction of G-diagrams with Kan extensions. We continue with enrichment and interactions with simplicial structure and finish the section with examples of G-diagrams. Section 2 mainly consists of the construction of a model structure on the category of G-diagrams in some nice model category. We also prove homotopy invariance and cofinality results for homotopy limits and colimits in this model category framework as well as an Elmendorf theorem. Section 3 deals with equivariant excision, and contains a conceptual proof of the Wirthmüller isomorphism, stated as a consequence of G-linearity for the identity functor on G-spectra. Finally, the appendix contains some technical results about G-diagrams indexed on directed Reedy categories and some properties of homotopy limits and colimits.

Acknowledgments

We wish to thank Irakli Pachkoria for help with proofreading and for pointing out a mistake in an earlier version of this article. We would also like to thank Ib Madsen for his enduring support and for encouraging us to write this paper.

1 Definitions and setup

1.1 Categories of G-diagrams

We first introduce some notation and conventions. If $\mathscr{C}$ is a (possibly large) category and I is a small category we write $\mathscr{C}^I$ for the usual category of functors from I to $\mathscr{C}$. By topological space we will mean compactly generated weak Hausdorff space and Top is the category of such spaces with continuous maps between them. We write $Map(X,Y)$ for the space of maps from X to Y endowed with the compact-open topology. The based variants of the above are Top_* and $Map_*(X,Y)$.

In the following $\mathscr{C}$ will be a category, G a finite group, and I a small category. By a slight abuse of notations we will also write G for the category with one object $*$ and one morphism $g\colon * \to *$ for each element $g \in G$, and with composition given by $g \circ h = gh$. The group G will act on I from the left and we will encode the action as a functor $a\colon G \to Cat$ sending $*$ to I. Most of the content of this section can be found in the work of Jackowski-Słomińska[JS01] or in Villarroel-Flores's paper [VF04].

Definition 1.1. (cf. [JS01, 2.2], [VF04, 3.1])Let $X\colon I \to \mathscr{C}$ be an I-shaped diagram in $\mathscr{C}$. A G-structure on X with respect to the action a is a collection of natural transformations $\{g_X\colon X \to X \circ a(g)\}$ such that

1. $e_X = id_X$
2. $(g_X)_{a(h)} \circ h_X = (gh)_X$ for all $g, h \in G$,

where $(g_X)_{a(h)}$ is the natural transformation obtained by restricting g_X along the functor $a(h)\colon I \to I$. An I-shaped diagram X with a G-structure will be called an I-shaped G-diagram in $\mathscr{C}$ with respect to the action a or simply a G-diagram in $\mathscr{C}$ if I and a are understood.

In order to simplify the notation we will mostly write g in stead of $a(g)$ when this does not cause confusion. Accordingly, when X and Y are I-indexed G-diagrams we will write f_g for the restriction of a map $f\colon X \to Y$ along the functor $g = a(g)\colon I \to I$. In the later sections we will sometimes write g instead of g_X.

Definition 1.2. A map of G-diagrams $f\colon X \to Y$ is a natural transformation $f\colon X \to Y$ of underlying diagrams such that for each $g \in G$ the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ {\scriptstyle g_X}\downarrow & & \downarrow{\scriptstyle g_Y} \\ X \circ g & \xrightarrow[f_g]{} & Y \circ g \end{array}$$

commutes in $\mathscr{C}^I$.

The composite of two maps of G-diagrams is again a map of G-diagrams. For a fixed action a of the group G on I we write $\mathscr{C}^I_a$ for the category whose objects are the G-diagrams in $\mathscr{C}$ with respect to a and with morphisms the maps of G-diagrams.

Let I and J be small categories with G-actions a and b respectively and let $F\colon I \to J$ be a functor. We say that F is G-equivariant if it commutes strictly with the G-actions, that is, if $F(gi) = gF(i)$ and $F(gf) = gF(f)$ for all objects i in I and morphisms f in I. If Y is a J-shaped G-diagram then the restriction $F^*Y = Y \circ F$ has a naturally induced G-structure with maps $g_{(F^*Y)} = F^*(g_Y)$.

Now assume that $\mathscr{C}$ is complete and cocomplete. Then the functor $F^*\colon \mathscr{C}^J \to \mathscr{C}^I$ has a left adjoint F_* and a right adjoint $F_!$ given by left and right Kan extension, respectively. We will now see that if X is an I-shaped G-diagram, then there are natural G-structures on F_*X and $F_!X$. We treat the left Kan extension first.

The value of the functor F_*X on an object j of J is given by the coequalizer

$$\coprod_{(i_0 \xrightarrow{\alpha} i_1,\, f\colon F(i_1)\to j)} X_{i_0} \underset{t}{\overset{s}{\rightrightarrows}} \coprod_{(i_0,\, f\colon F(i_0)\to j)} X_{i_0} \twoheadrightarrow F_*X_j,$$

where s projects onto the source of the indexing map α and t maps into the target of α by the map $X(\alpha)$. For an element $g \in G$ the natural transformation g_X induces a map of diagrams

$$\begin{array}{ccccc} \coprod_{(i_0 \xrightarrow{\alpha} i_1,\, f\colon F(i_1)\to j)} X_{i_0} & \underset{t}{\overset{s}{\rightrightarrows}} & \coprod_{(i_0,\, f\colon F(i_0)\to j)} X_{i_0} & \twoheadrightarrow & F_*X_j \\ \downarrow{\scriptstyle \coprod g_{X_{i_0}}} & & \downarrow{\scriptstyle \coprod g_{X_{i_0}}} & & \vdots\ {\scriptstyle g_{F_*X_j}} \\ \coprod_{(i'_0 \xrightarrow{\alpha'} i'_1,\, f'\colon F(i'_1)\to gj)} X_{i'_0} & \underset{t}{\overset{s}{\rightrightarrows}} & \coprod_{(i'_0,\, f'\colon F(i'_0)\to gj)} X_{i'_0} & \twoheadrightarrow & F_*X_{gj} \end{array}$$

and the dotted arrow is the j-component of the natural transformation $g_{F_*X}\colon F_*X \to (F_*X) \circ g$. It is not hard to see that the set $\{g_{F_*X}\}_{g\in G}$ constitutes a G-structure on F_*X and that the underlying functor F_* takes maps of I-indexed G-diagrams to maps of J-indexed G-diagrams. Similarly, for the right Kan extension $F_!$ a dual construction with equalizers gives a G-structure $\{g_{F_!X}\}_{g\in G}$ on $F_!X$. We write simply F_*X and $F_!X$ for the G-diagrams obtained in this way.

Proposition 1.3: *The constructions F_*X and $F_!X$ define functors $F_*\colon \mathscr{C}^I_a \to \mathscr{C}^J_b$ and $F_!\colon \mathscr{C}^I_a \to \mathscr{C}^J_b$.*

A particularly interesting case of the above is when $J = *$ the category with one object and one morphism and trivial G-action. In this case the functors F_* and $F_!$ are more commonly known as colim_I and $\lim_I$, respectively.

Corollary 1.4: *Let X be an I-indexed G-diagram. Then the above constructions induce natural left G-actions on* $\operatorname{colim}_I X$ *and* $\lim_I X$.

Example 1.5. (Products and coproducts) Let I be a discrete category with G-action, i.e., a G-set and consider a G-diagram X in the category Set of sets. The coproduct $\coprod_I X$ is the set of pairs (i, x) where $x \in X_i$ and the action of $g \in G$ is given by

$$g(x, i) = (g_{X_i}(x), gi).$$

The product $\prod_I X$ is the set of functions $\mathbf{x}\colon I \to \bigcup_{i\in I} X_i$ such that $\mathbf{x}(i) \in X_i$ for all $i \in I$. The action of $g \in G$ on $\mathbf{x} \in \prod_I X$ is determined by the equation

$$(g\mathbf{x})(gi) = g_{X_i}(\mathbf{x}(i)).$$

This example generalizes to arbitrary categories with products and coproducts but the notation becomes more cumbersome when one can no longer speak about elements of objects.

We now give an alternative description of G-diagrams which is sometimes easier to work with.

Definition 1.6. The category $G \rtimes_a I$ is given as follows:

- $obG \rtimes_a I = obI$
- A morphism $i \to j$ in $G \rtimes_a I$ is a pair $(g, f\colon gi \to j)$ where $g \in G$.
- Composition is given by $(h, f'\colon hj \to k) \circ (g, f\colon gi \to j) = (gh, f' \circ hf\colon ghi \to k)$.

Remark 1.7. Note that in the notation of e.g. [Tho79] this is the Grothendieck construction $G \int a$.

A G-diagram X gives rise to a functor $X^{\rtimes_a}\colon G \rtimes_a I \to \mathscr{C}$ by setting

$$X_i^{\rtimes_a} = X_i$$

on objects, and by setting

$$X^{\rtimes_a}(g, f\colon gi \to j) = X(f) \circ g_{X_i}$$

on morphisms. We leave it to the reader to check that this respects composition of maps.

Lemma 1.8: *The assignment $X \mapsto X^{\rtimes_a}$ is functorial and defines an isomorphism of categories*

$$\Phi\colon \mathscr{C}_a^I \xrightarrow{\cong} \mathscr{C}^{G\rtimes_a I}.$$

Proof. The functoriality is clear. We will define a functor $\Phi'\colon \mathscr{C}^{G\rtimes_a I} \to \mathscr{C}_a^I$ which is inverse to Φ.

For a diagram $Y\colon G \rtimes_a I \to \mathscr{C}$ define the underlying diagram of $\Phi'(Y)$ to be $(Y|_I)$, i.e., the restriction of Y along the canonical inclusion $\iota\colon I \hookrightarrow G \rtimes_a I$ given by $\iota(i) = i$ and $\iota(f\colon i \to j) = (e, f\colon i \to j)$. For an element $g \in G$ the natural transformation g_Y is defined by $g_{Y_i} = Y(g, id\colon gi \to gi)$. Both naturality of the g_Y's and conditions 1) and 2) of Definition 1.1 follow from the functoriality of Y with respect to morphisms in $G \rtimes_a I$. For a natural transformation $\alpha\colon Y \to Z$ in $\mathscr{C}^{G\rtimes_a I}$ we define $\Phi'(\alpha) = \alpha|_I$. It is now easy to check that the functors Φ and Φ' are mutually inverse. □

Corollary 1.9: *Let $\mathscr{C}$ be a bicomplete category. Then $\mathscr{C}_a^I$ is also bicomplete.*

Proof. The diagram category $\mathscr{C}^{G\rtimes_a I}$ is bicomplete since $\mathscr{C}$ is. It follows that $\mathscr{C}_a^I$ is bicomplete. □

1.2 Enrichments and homotopy (co)limits

If $\mathscr{C}$ is any category, then the category $\mathscr{C}^G$ is naturally enriched in left G-sets in the following way. For objects c, d of $\mathscr{C}^G$ let $\mathscr{C}(c, d)$ be the set of maps between the underlying objects in $\mathscr{C}$. Then G acts on $\mathscr{C}(c, d)$ by conjugation

$$g \cdot f = g_d \circ f \circ (g^{-1})_c$$

where $(g^{-1})_c$ and g_d represent the actions of g^{-1} and g on c and d respectively. The fixed points $\mathscr{C}(c,d)^G$ are precisely the G-equivariant maps from c to d.

If I is small a category with an action a of G, then the category $\mathscr{C}_a^I$ of G-diagrams becomes enriched in left G-sets by taking $\underline{\mathscr{C}}_a^I(X,Y)$ to be the set $\mathscr{C}^I(X,Y)$ of maps of underlying diagrams $f\colon X \to Y$ with action given by

$$g \cdot f = (g_Y)_{g^{-1}} \circ f_{g^{-1}} \circ (g^{-1})_X.$$

If f is fixed under the action of G, then

$$f = g^{-1}f = ((g^{-1})_Y)_g \circ f_g \circ g_X = (g_Y)^{-1} \circ f_g \circ g_X.$$

In other words, f is fixed if and only if the square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ {\scriptstyle g_X}\downarrow & & \downarrow{\scriptstyle g_Y} \\ X \circ g & \xrightarrow[f_g]{} & Y \circ g \end{array}$$

commutes for all $g \in G$. It follows that the fixed points $\underline{\mathscr{C}}_a^I(X,Y)^G$ are precisely the maps of G-diagrams $\mathscr{C}_a^I(X,Y)$. If $I = *$ then this statement reduces to the one above about maps in $\mathscr{C}^G$.

Proposition 1.10: *Let I and J be small categories with G-actions a and b, respectively and let $F\colon I \to J$ be an equivariant functor. Then, for X an I-indexed G-diagram and Y a J-indexed G-diagram the bijections*

$$\alpha_{X,Y}\colon \underline{\mathscr{C}}_a^I(X, F^*Y) \xrightarrow{\cong} \underline{\mathscr{C}}_b^J(F_*X, Y)$$

and

$$\beta_{X,Y}\colon \underline{\mathscr{C}}_a^I(F^*X, Y) \xrightarrow{\cong} \underline{\mathscr{C}}_b^J(X, F_!Y)$$

induced by the adjunctions on underlying diagrams are G-equivariant.

Proof. We show that $\alpha = \alpha_{X,Y}$ is equivariant, the argument for $\beta_{X,Y}$ is similar.

Let $f\colon X \to F^*Y$ be a map of diagrams and $g \in G$. Then $\alpha(g \cdot f)$ is the unique map $F_*X \to Y$ such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{g \cdot f} & F^*Y \\ {\scriptstyle \eta_X}\downarrow & \nearrow_{F^*(\alpha(g\cdot f))} & \\ F^*F_*X & & \end{array} \tag{2}$$

commutes, where η_X is the unit of the (F_*, F^*)-adjunction at the object X. Consider the following diagram:

$$\begin{array}{ccccccc} X & \xrightarrow{(g^{-1})_X} & X \circ g^{-1} & \xrightarrow{f_{g^{-1}}} & (F^*Y) \circ g^{-1} & \xrightarrow{(F^*g_Y)_{g^{-1}}} & F^*Y \\ {\scriptstyle \eta_X}\downarrow & & {\scriptstyle \eta_{X,g^{-1}}}\downarrow & & \downarrow{=} & \nearrow_{(F^*g_Y)_{g^{-1}}} & \\ F^*F_*X & \xrightarrow[F^*((g^{-1})_{F_*X})]{} & (F^*F_*X) \circ g^{-1} & \xrightarrow[F^*\alpha(f)_{g^{-1}}]{} & (F^*Y) \circ g^{-1}. & & \end{array}$$

The commutativity of the left hand square follows immediately from the definition of g_{F_*X} and middle square commutes by the definition of $\alpha(f)$. Composing the maps in the top row gives $(F^*g_Y)_{g^{-1}} \circ f_{g^{-1}} \circ (g^{-1})_X = g \cdot f$ and composing along the bottom row from F^*F_*X to F^*Y gives

$$F^*((g_Y)_{g^{-1}} \circ \alpha(f)_{g^{-1}} \circ (g^{-1})_{F_*X}) = F^*(g \cdot \alpha(f)).$$

It follows that $F^*(g \cdot \alpha(f))$ defines a lift in the diagram (2) so, by uniqueness of the lift, we conclude that $\alpha(g \cdot f) = g \cdot \alpha(f)$. □

Taking fixed points in Proposition 1.10 we immediately get the following:

Corollary 1.11: *The functors F_* and $F_!$ are left and right adjoint, respectively, to the restriction functor $F^*\colon \mathscr{C}_b^J \to \mathscr{C}_a^I$. In particular the diagonal $\Delta_I\colon \mathscr{C}^G \to \mathscr{C}_a^I$ has left and right adjoints colim_I and $\lim_I$, respectively.*

Let I be a category with G-action a and let G act diagonally on the product $I^{op} \times I$. Given a G-diagram $Z\colon I^{op} \times I \to \mathscr{C}$ recall that the *end* $\int_i Z_{i,i}$ of Z is the equalizer

$$\int_i Z_{i,i} \rightarrowtail \prod_i Z_{i,i} \underset{t}{\overset{s}{\rightrightarrows}} \prod_{f\colon i\to j} Z_{j,i}$$

where s and t act on the left and right, respectively by the map f. The end $\int_i Z_{i,i}$ inherits a left G-action by the maps

$$\begin{array}{ccccc} \int_i Z_{i,i} & \longrightarrow & \prod_i Z_{i,i} & \underset{t}{\overset{s}{\rightrightarrows}} & \prod_{f\colon i\to j} Z_{j,i} \\ \downarrow{\scriptstyle g_{(\int Z)}} & & \downarrow{\scriptstyle \prod_i g_{Z_{(i,i)}}} & & \downarrow{\scriptstyle \prod_f g_{Z_{(j,i)}}} \\ \int_i Z_{i,i} & \longrightarrow & \prod_i Z_{i,i} & \underset{t}{\overset{s}{\rightrightarrows}} & \prod_{f\colon i\to j} Z_{j,i} \end{array} \qquad (*)$$

The *coend* $\int^i Z_{i,i}$ is the coequalizer

$$\coprod_{f\colon i\to j} Z_{j,i} \underset{t}{\overset{s}{\rightrightarrows}} \coprod_i Z_{i,i} \twoheadrightarrow \int_i Z_{i,i}.$$

which inherits a G-action in a similar way.

Example 1.12. If $X, Y\colon I \to \mathscr{C}$ are diagrams in $\mathscr{C}$ then we can describe the set of maps (natural transformations) between them as the end

$$\mathscr{C}^I(X,Y) = \int_i \mathscr{C}(X_i, Y_i).$$

Similarly, for G-diagrams X, Y in $\mathscr{C}_a^I$ there is a natural isomorphism of G-sets

$$\underline{\mathscr{C}}_a^I(X,Y) \cong \int_i \mathscr{C}(X_i, Y_i)$$

with the G-action on the left hand as described above.

By a *simplicial category* we will mean a category $\mathscr{C}$ that is enriched, tensored and cotensored in simplicial sets, in the sense of e.g. [DS07, 2.2] or [GJ09, II,2.1]. This means that for any two objects c and d in $\mathscr{C}$ there is a simplicial set $Map_{\mathscr{C}}(c,d)$, and a natural bijection $\mathscr{C}(c,d) \cong Map_{\mathscr{C}}(c,d)_0$. Moreover, given a simplicial set K there are objects $K \otimes c$ and $map_{\mathscr{C}}(K,c)$ of $\mathscr{C}$. These satisfy some associativity constraints and naturality conditions making $Map_{\mathscr{C}}(-,-)$ and $map_{\mathscr{C}}(-,-)$ contravariant functors in the first variable and covariant in the second variable and $-\otimes-$ covariant in both variables. Finally for all c, d in $\mathscr{C}$ and K in $sSet$ there are natural isomorphisms in $sSet$

$$Map_{\mathscr{C}}(c \otimes K, d) \cong Map(K, Map_{\mathscr{C}}(c,d)) \cong Map_{\mathscr{C}}(c, map_{\mathscr{C}}(K,d)),$$

where Map with no subscript denotes the usual internal hom-object in $sSet$.

Using this structure we will now describe additional structure on the category $\mathscr{C}_a^I$ of I-indexed G-diagrams in a simplicial category $\mathscr{C}$. We begin with enrichment. We noted above that for a pair X, Y of G-diagrams in $\mathscr{C}$ the set $\mathscr{C}^I(X,Y)$ has a G-action induced by the G-structures on X and Y. This gives $\mathscr{C}_a^I$ the structure of a category enriched in left G-sets. The functor $i,j \mapsto Map_{\mathscr{C}}(X_i, Y_j)$ going from $I^{op} \times I$ to $sSet$ becomes a G-diagram by letting $g \in G$ act at i,j by

$$Map_{\mathscr{C}}(g_{X_i}^{-1}, g_{Y_j})\colon Map_{\mathscr{C}}(X_i, Y_j) \to Map_{\mathscr{C}}(X_{gi}, Y_{gj}).$$

Definition 1.13. With X, Y as above, set

$$Map_{\mathscr{C}_a^I}(X,Y) = \int_i Map_{\mathscr{C}}(X_i, Y_i)$$

with the G-action as described in the diagram $(*)$.

In other words the mapping space $Map_{\mathscr{C}_a^I}(X,Y)$ is the equalizer

$$Map_{\mathscr{C}_a^I}(X,Y) \longrightarrow \prod_i Map_{\mathscr{C}}(X_i, Y_i) \underset{t}{\overset{s}{\rightrightarrows}} \prod_{f\colon i \to j} Map_{\mathscr{C}}(X_j, Y_i)$$

It is not hard to see that this defines an enrichment of $\mathscr{C}_a^I$ in $sSet^G$ and that for each $n \geq 0$ there is an isomorphism of G-sets

$$Map_{\mathscr{C}_a^I}(X,Y)_n \cong \underline{\mathscr{C}}_a^I(\Delta^n \otimes X, Y).$$

Definition 1.14. Let $K\colon I \to sSet$, $L\colon I^{op} \to sSet$, and $X\colon I \to \mathscr{C}$ be G-diagrams. We set

$$map_I^a(K,X) = \int_i map_{\mathscr{C}}(K_i, X_i) \tag{3}$$

$$L \otimes_I^a X = \int^i L_i \otimes X_i \tag{4}$$

and give both the G-actions as in $(*)$.

Let $[n]$ be the usual category with objects $0, 1 \ldots n$ and a morphism $i \to j$ if and only if $i \leq j$. For a small category I the nerve NI is the usual simplicial set with $NI_n = Fun([n], I)$. Taking over-categories gives a functor $N(I/-)\colon I \to sSet$. The G-action on I gives maps $N_{/i,g}\colon N(I/i) \to N(I/gi)$ for $g \in G$ and i an object of I. These maps combine to give a G-diagram structure on $N(I/-)$. Similarly the functor $N(-/I)^{op}\colon I^{op} \to sSet$ with the maps $N_{i,g/}\colon N(i/I)^{op} \to N(gi/I)^{op}$ defines a G-diagram in $sSet$.

Definition 1.15. For a G-diagram X in $\mathscr{C}$ the homotopy limit of X is

$$\operatorname*{holim}_I X = map_I^a(N(I/-), X)$$

and the homotopy colimit of X is

$$\operatorname*{hocolim}_I X = N(-/I)^{op} \otimes_I^a X.$$

In the presence of a model structure the words homotopy limit and colimit will always refer to these particular construction and not, a priori, the derived functors of the limit and colimit respectively.

Note that there are maps of diagrams $N(-/I)^{op} \to *$ and $N(I/-) \to *$, where $*$ denotes a chosen one-point simplicial set in both cases. From the formulas above it is easy to see that there are natural

isomorphisms $map^a_I(*, X) \cong \lim X$ and $X \otimes^a_I * \cong \operatorname{colim} X$. The maps to the terminal diagrams induce maps

$$\lim X \to \operatorname{holim} X$$

and

$$\operatorname{hocolim} X \to \operatorname{colim} X.$$

This paper is in part motivated by the question "when are these maps weak equivalences in $\mathscr{C}^G$?"

1.3 Examples of G-diagrams

In this section we will provide many of the motivating examples for the theory of G-diagrams. The diagrams will usually have values in the category Top_* of pointed spaces.

For the first two examples we need to fix some notation. Let Z be a pointed space with an action by the finite group G. If T is a finite left G-set then we write $\mathbb{R}[T]$ for the permutation representation with basis $\{e_t\}_{t\in T}$. The subspace of $\mathbb{R}[T]$ generated by the element $N_T = \sum_{t\in T} e_t$ is a one-dimensional trivial subrepresentation of $\mathbb{R}[T]$. We define $S^{\tilde{T}}$ to be the one-point compactification of the orthogonal complement of $\mathbb{R} \cdot N_T$ under the usual inner product. We write $\Omega^{\tilde{T}} Z$ for the G-space of continuous pointed maps $Map_*(S^{\tilde{T}}, Z)$ with the conjugation action of G and $\Sigma^{\tilde{T}} Z$ for the smash product $S^{\tilde{T}} \wedge Z$ with the diagonal G-action.

Example 1.16. The power set $\mathcal{P}(T)$ inherits a left G-action from the action on T. We think of the poset $\mathcal{P}(T) \setminus \emptyset$ as a category with G-action. Let $\omega^{\tilde{T}} Z$ be the $\mathcal{P}(T) \setminus \emptyset$-indexed G-diagram whose value on a subset $U \subseteq T$ is $*$ if $U \neq T$ and Z if $U = T$. The G-structure on $\omega^{\tilde{T}} Z$ is given by the action of G on Z at the fixed object T and by the unique maps $* \to *$ elsewhere in the diagram. We claim that there is a G-homeomorphism

$$\operatorname*{holim}_{\mathcal{P}(T)\setminus\emptyset} \omega^{\tilde{T}} Z \cong \Omega^{\tilde{T}} Z$$

which is natural in Z. To see this we begin by noting that $|N((\mathcal{P}(T) \setminus \emptyset)/U)| \cong |N(\mathcal{P}(U) \setminus \emptyset)|$ is G-homeomorphic to the (barycentric subdivision of the) standard simplex Δ^U in $\mathbb{R}[U]$ spanned by all the basis vectors. Since $\omega^{\tilde{T}} Z$ has all entries trivial except at the last vertex T we see that $\operatorname{holim} \omega^{\tilde{T}} Z$ is homeomorphic to the subspace in $Map(\Delta^U, Z)$ of maps whose restriction to the boundary is the constant map to the base-point of Z. The naturality is clear, so this proves the claim.

Example 1.17. Similarly, we think of the poset $\mathcal{P}(T) \setminus T$ as a category with G-action and define the G-diagram $\sigma^{\tilde{T}} Z$ to have the value Z at the vertex $\emptyset$ and $*$ elsewhere. The G-diagram structure is induced by the G-action on Z and the unique maps $* \to *$. A similar argument to the one for $\omega^{\tilde{T}} Z$ shows that there is a natural G-homeomorphism

$$\operatorname*{hocolim}_{\mathcal{P}(T)\setminus T} \sigma^{\tilde{T}} Z \cong \Sigma^{\tilde{T}} Z.$$

Example 1.18. More generally, for any pointed object c of $\mathscr{C}^G$ define the $\tilde{T}$-loop space and $\tilde{T}$-suspension of c as the pullback and pushout in $\mathscr{C}^G$

$$\begin{array}{ccc} \Omega^{\tilde{T}} c & \longrightarrow & map_{\mathscr{C}}(N\mathcal{P}(T)\setminus\emptyset, c) \\ \downarrow & & \downarrow \\ * & \longrightarrow & map_{\mathscr{C}}(\partial N\mathcal{P}(T)\setminus\emptyset, c), \end{array} \qquad \begin{array}{ccc} (\partial N\mathcal{P}(T)^{op}\setminus T) \otimes c & \longrightarrow & * \\ \downarrow & & \downarrow \\ (N\mathcal{P}(T)^{op}\setminus T) \otimes c & \longrightarrow & \Sigma^{\tilde{T}} c. \end{array}$$

In the case of a pointed space we recover the usual loop and suspension spaces (the second construction does not require a base-point, and in the case of an unpointed space it gives the unreduced suspension).

As in the previous examples there are natural isomorphisms in $\mathscr{C}^G$

$$\operatorname*{holim}_{\mathcal{P}(T)\setminus\emptyset} \omega^{\tilde{T}} c \cong \Omega^{\tilde{T}} c \qquad \operatorname*{hocolim}_{\mathcal{P}(T)\setminus T} \sigma^{\tilde{T}} c \cong \Sigma^{\tilde{T}} c$$

Example 1.19. We already saw that for a category I with G-action the functor $N(I/-)$ to $sSet$ has an obvious G-structure. For a functor $F\colon I \to J$ and an object j of J one can form the over-category F/j and the assignment $j \mapsto N(F/j)$ defines a functor $N(F/-)\colon J \to sSet$. If F is an equivariant functor between categories with G-action there are functors $F/j \to F/(gj)$ induced by the G-actions and after applying the nerve these give a G-structure on the diagram $N(F/-)$. In fact, $N(F/-)$ with this G-structure is the left Ken extension $F_*N(I/-)$ of $N(I/-)$ along F. This will be important later when we discuss homotopy cofinality and cofibrancy of G-diagrams.

Example 1.20. Let $X\colon I \to \mathscr{C}$ be a diagram in a simplicial category $\mathscr{C}$. Define the diagram qX by $qX_i = \operatorname{hocolim}_{I/i} u_i^* X$ where $u_i\colon I/i \to I$ is the functor that forgets the map to i. A map $f\colon i \to j$ in I induces a functor $I/i \to I/j$ and hence a map $qX_i \to qX_j$. The natural map from the homotopy colimit to the colimit induces maps

$$qX_i = \operatorname*{hocolim}_{I/i} u_i^* X \to \operatorname*{colim}_{I/i} u_i^* X \xrightarrow{\cong} X_i,$$

which combine to a map of diagrams $\rho_X\colon qX \to X$. If X is a G-diagram then the functor $I/i \to I/gi$ induced by multiplication by $g \in G$ induces a map $qX_i \to qX_{gi}$ and together these maps constitute a G-structure on qX. It is a classical fact that the objects $\operatorname{colim}_I qX$ and $\operatorname{hocolim}_I X$ are isomorphic and in ii of Proposition 2.16 we prove that this isomorphism is G-equivariant when X is a G-diagram.

2 G-diagrams and model structures

2.1 Equivariant model categories

Let $\mathscr{C}$ be a complete and cocomplete category, G a finite group and $H, H' \leq G$ a pair of subgroups. A finite set K with commuting left H'-action and right H-action induces a pair of adjoint functors

$$K \otimes_H (-)\colon \mathscr{C}^H \rightleftarrows \mathscr{C}^{H'}\colon \hom_{H'}(K, -)$$

The left adjoint is defined as

$$K \otimes_H c = \operatorname{colim}\left(H \xrightarrow{\coprod_K c} \mathscr{C}\right)$$

where $\coprod_K c$ is the H-colimit of the constant H-diagram Δc on the discrete H-category K^δ, (see Example 1.5), and the H'-action is induced by the H'-action on K. Dually, define

$$\hom_{H'}(K, d) = \lim\left(H' \xrightarrow{\prod_K c} \mathscr{C}\right)$$

with left H-action defined by right action on K. These functors are adjoint by the sequence of natural isomorphisms

$$\begin{aligned}&\mathscr{C}^{H'}(K \otimes_H c, d) \cong \mathscr{C}(K \otimes_H c, d)^{H'} \cong \mathscr{C}^H(\textstyle\coprod_K c, d)^{H'} \cong \\ &\mathscr{C}^K_a(\Delta_K c, \Delta_K d)^{H'} \cong \mathscr{C}^H(c, \textstyle\prod_K d)^{H'} \cong \mathscr{C}^H(c, \lim_{H'}(\prod_K d)) = \\ &\mathscr{C}^H(c, \hom_{H'}(K, d))\end{aligned}$$

In the following we will always use the fixed point model structure on $sSet^G$ (see e.g. [Shi03, 1.2]) unless otherwise is stated.

Definition 2.1. A G-model category is a cofibrantly generated simplicial model category $\mathscr{C}$, together with the data of a cofibrantly generated model structure on $\mathscr{C}^H$ for every subgroup $H \leq G$, satisfying

1. The model structure on $\mathscr{C}^H$ together with the $sSet^H$-enrichment, tensored and cotensored structures induced from $\mathscr{C}$ forms a cofibrantly generated $sSet^H$-enriched model structure on $\mathscr{C}^H$,
2. For every pair of subgroups $H, H' \leq G$, and finite set K with commuting free left H'-action and free right H-action the adjunction

$$K \otimes_H (-) \colon \mathscr{C}^H \rightleftarrows \mathscr{C}^{H'} \colon \hom_{H'}(K, -)$$

is a Quillen adjunction.

Remark 2.2. For $H' \leq H$ and $K = H$ with actions given by left H' and right H multiplications, the functor

$$H \otimes_H (-) \colon \mathscr{C}^H \longrightarrow \mathscr{C}^{H'}$$

is isomorphic to the functor $\mathrm{res}^H_{H'}$ that restricts the action. Similarly for $K = H$ with left H multiplication and right H' multiplication the functor

$$\hom_H(H, -) \colon \mathscr{C}^H \longrightarrow \mathscr{C}^{H'}$$

is also isomorphic to the functor $\mathrm{res}^H_{H'}$. It follows from the second condition that $\mathrm{res}^H_{H'}$ is both a left and a right Quillen functor, and therefore it preserves cofibrations, acyclic cofibrations, fibrations, acyclic fibrations and equivalences between cofibrant or fibrant objects.

Example 2.3. Let $\mathscr{C}$ be a cofibrantly generated $sSet$-enriched model category. The collection of projective model structures (naive) on $\mathscr{C}^H$ for $H \leq G$ defines a G-model structure on $\mathscr{C}^G$. To see this, just notice that since the H'-action on K is free, a choice of section for the quotient map $K \to H' \backslash K$ induces a natural isomorphism

$$\mathrm{res}^H_e \hom_{H'}(K, c) \cong \prod_{H' \backslash K} c$$

where $\mathrm{res}^H_e : \mathscr{C}^H \to \mathscr{C}$ is the forgetful functor. Therefore $\hom_{H'}(K, -)$ preserves fibrations and acyclic fibrations.

Example 2.4. Let $\mathscr{C}$ be a cofibrantly generated $sSet$-enriched model category, and fix a pair of finite groups $H \leq G$. For all subgroup $L \leq H$, the L-fixed points functor $(-)^L \colon \mathscr{C}^H \to \mathscr{C}$ is defined as the composite

$$\mathscr{C}^H \xrightarrow{U} \mathscr{C}^L \xrightarrow{\lim} \mathscr{C}$$

If these functors are cellular in the sense of [GM13], the category $\mathscr{C}^H$ inherits a $sSet^H$-enriched model structure where weak equivalences and fibrations are the maps that are sent by every functor $(-)^L$ to weak equivalences and fibrations of $\mathscr{C}$ respectively (cf. [MM02, 2.8],[GM13],[Ste10]). In particular this construction gives the standard model structure on (pointed) spaces with H-action.

The collection of the model categories $\mathscr{C}^H$, for H running over the subgroups subgroups of G, assemble into a G-model category. Let us see that the left adjoint $K \otimes_H (-)$ is a left Quillen functor. The generating cofibrations of $\mathscr{C}^H$ are by definition the images of the generating cofibrations of $\mathscr{C}$ by the functors

$$J \otimes (-) \colon \mathscr{C} \longrightarrow \mathscr{C}^H$$

where J ranges over finite sets with left H-action. Similarly for generating acyclic cofibrations. There is a natural isomorphism

$$K \otimes_H (J \otimes (-)) \cong (K \times_H J) \otimes (-)$$

and the right hand functor preserves cofibrations and acyclic cofibrations by assumption. Thus $K \otimes_H (-)$ preserves generating (acyclic) cofibrations. Since it is a left adjoint it preserves colimits, and therefore all (acyclic) cofibrations (see e.g. [Hir03, 11.2]).

Example 2.5. Let $\mathscr{C} = \mathrm{Sp}^O$ be the category of orthogonal spectra and G a finite group. The category $(\mathrm{Sp}^O)^G$ of G-objects in Sp^O is naturally equivalent to the category of orthogonal G-spectra $\mathscr{I}_G^{\mathscr{V}}\mathscr{S}$ of [MM02] indexed on a universe $\mathscr{V}$ for finite dimensional G-representations (cf.[MM02, V.1], [Sch13, 2.7]). Given any subgroup $H \leq G$, we endow $(\mathrm{Sp}^O)^H$ with the model structure induced by the stable model structure on $\mathscr{I}_H^{i^*\mathscr{V}}\mathscr{S}$ of [MM02] under the equivalence of categories $(\mathrm{Sp}^O)^H \simeq \mathscr{I}_H^{i^*\mathscr{V}}\mathscr{S}$. Here $i\colon H \to G$ denotes the inclusion, and $i^*\mathscr{V}$ is the universe of representations of H that are restrictions of representations of G in $\mathscr{V}$. The adjunctions

$$K \otimes_H (-)\colon (\mathrm{Sp}^O)^H \rightleftarrows (\mathrm{Sp}^O)^{H'}\colon \hom_{H'}(K, -)$$

are the standard induction-coinduction adjunctions, and they are Quillen adjunctions by [MM02, V-2.3]. The collection of model categories $\{(\mathrm{Sp}^O)^H\}_{H\leq G}$ forms a G-model category.

2.2 A model structure on G-diagrams

Let G be a finite group, $\mathscr{C}$ a category, and I a small category with G-action a. Given a G-diagram X in $\mathscr{C}_a^I$ and an object $i \in I$, the vertex $X_i \in \mathscr{C}$ inherits an action by the stabilizer group $G_i \leq G$ of the object i. This gives an evaluation functor $\mathrm{ev}_i\colon \mathscr{C}_a^I \to \mathscr{C}^{G_i}$ for every object i.

Theorem 2.6: *Let $\mathscr{C}$ be a G-model category (see 2.1). There is a cofibrantly generated $sSet^G$-enriched model structure on the category of G-diagrams $\mathscr{C}_a^I$ with*

1. *weak equivalences the maps of G-diagrams $f\colon X \to Y$ whose restrictions $\mathrm{ev}_i f$ are weak equivalences in $\mathscr{C}^{G_i}$ for every $i \in I$,*
2. *fibrations the maps of G-diagrams $f\colon X \to Y$ whose restrictions $\mathrm{ev}_i f$ are fibrations in $\mathscr{C}^{G_i}$ for every $i \in I$,*
3. *generating cofibrations and acyclic cofibrations*

$$F\mathcal{I} = \bigcup_{i\in I} F_i\mathcal{I}_i \qquad \text{and} \qquad F\mathcal{J} = \bigcup_{i\in I} F_i\mathcal{J}_i$$

where $\mathcal{I}_i$ and $\mathcal{J}_i$ are respectively generating cofibrations and acyclic cofibrations of $\mathscr{C}^{G_i}$, and $F_i\colon \mathscr{C}^{G_i} \to \mathscr{C}_a^I$ is the left adjoint to the evaluation functor ev_i.

Remark 2.7. Under the isomorphism $\mathscr{C}_a^I \cong \mathscr{C}^{I\ltimes G}$ of Lemma 1.8 the evaluation functor ev_i corresponds to restriction along the functor $\iota_i\colon G_i \to I \ltimes G$ that sends the unique object to i and a morphism g to $(g, \mathrm{id}_i\colon gi = i \to i)$. Since $\mathscr{C}$ has all colimits a left adjoint for ev_i exists.

Before proving the theorem we need to identify the left adjoints of the evaluation functors. For fixed objects $i, j \in I$ let K_{ji} be the set of morphisms

$$K_{ji} = \hom_{I\ltimes G}(i, j) = \{(g \in G, \alpha\colon gi \to j)\}$$

The stabilizer group G_j acts freely on the left on K_{ji} by left multiplication on G and by the category action on the morphism component. The group G_i acts freely on the right on K_{ji} by right multiplication on the G-component.

For every $c \in \mathscr{C}^{G_i}$ define a diagram $F_ic\colon I \to C$ by sending an object $j \in I$ to

$$(F_ic)_j = K_{ji} \otimes_{G_i} c$$

A morphism $\beta\colon j \to j'$ in I induces a map $(F_ic)_j \to (F_ic)_{j'}$ via the G_i-equivariant map $\beta_*\colon K_{ji} \to K_{j'i}$

$$\beta_*(g, \alpha\colon gi \to j) = (g, \beta \circ \alpha)$$

The G_i-equivariant maps $g\colon K_{ji} \to K_{(gj)i}$

$$g(g', \alpha\colon g'i \to j) = (gg', g\alpha\colon gg'i \to gj)$$

define a G-structure on F_ic.

Lemma 2.8: *The functor $F_i\colon \mathscr{C}^{G_i} \to \mathscr{C}_a^I$ is left adjoint to the evaluation functor $\mathrm{ev}_i\colon \mathscr{C}_a^I \to \mathscr{C}^{G_i}$.*

Proof. We prove that under the isomorphism $\mathscr{C}_a^I \cong \mathscr{C}^{I \ltimes G}$ of Lemma 1.8 the functor F_i corresponds to the left Kan extension along the inclusion $\iota_i\colon G_i \to I \ltimes G$. For an object $j \in I$, the category ι_i/j is the disjoint union of categories

$$\iota_i/j = \coprod_{\substack{z \in G/G_i \\ zi \to j}} Ez$$

where Ez is the translation category of the right G_i-set z, with one object for every element of the orbit z, and a unique morphism $h\colon g \to g'$ whenever $g' = gh^{-1}$ for some $h \in G_i$. An object $c \in \mathscr{C}^{G_i}$ induces a diagram $Ec\colon Ez \to G_i \xrightarrow{c} \mathscr{C}$, where the first functor collapses all the objects to the unique object of G_i, and sends the unique morphism $g \to gh^{-1}$ to h. The left Kan extension along ι_i at c is by definition the diagram $L_i c$ with j-vertex

$$(L_i c)_j = \coprod_{\substack{z \in G/G_i \\ zi \to j}} \operatorname*{colim}_{Ez} Ec$$

Notice that the indexing set of the coproduct is precisely the orbit set K_{ji}/G_i. There is a canonical map of diagrams $F_i c \to L_j c$, which at a vertex j is induced by

$$\coprod_{K_{ji}} c \longrightarrow \coprod_{K_{ji}/G_i} \operatorname*{colim}_{Ez} Ec = (L_i c)_j$$

which on the (g,α)-component is the canonical map $c = (Ec)_g \to \mathrm{colim}_{E[g]} Ec$ to the $[g,\alpha]$-coproduct component. This map respects the G_j-structure, which on $L_i c$ acts via the indexing set K_{ji}/G_i. To show that it is an isomorphism, choose a section $s\colon G/G_i \to G$ for the projection map. This gives a map

$$(L_i c)_j = \coprod_{K_{ji}/G_i} \operatorname*{colim}_{Ez} Ec \longrightarrow \coprod_{K_{ji}} c \longrightarrow K_{ji} \otimes_{G_i} c = (F_i c)_j$$

that on the (z,α)-component is the map induced by $s(z)^{-1}g\colon (Ez)_g = c \to c$ to the $(s(z),\alpha)$-component. □

Proof of 2.6. Weak equivalences and fibrations in $\mathscr{C}_a^I$ are by definition the morphisms that are sent to weak equivalences and fibrations, respectively, by the functor

$$\prod_{i \in I} \mathrm{ev}_i\colon \mathscr{C}_a^I \longrightarrow \prod_{i \in I} \mathscr{C}^{G_i}$$

It follows from Lemma 2.8 that the coproduct of the functors F_i defines a left adjoint

$$L\colon \prod_{i \in I} \mathscr{C}^{G_i} \xrightarrow{\prod F_i} \prod_{i \in I} \mathscr{C}_a^I \xrightarrow{\coprod} \mathscr{C}_a^I$$

for the product of the evaluation functors. The collections

$$\mathcal{I} = \bigcup_{i \in I} (\mathcal{I}_i \times \prod_{j \neq i} \mathrm{id}_{\emptyset_j}) \quad \text{and} \quad \mathcal{J} = \bigcup_{i \in I} (\mathcal{J}_i \times \prod_{j \neq i} \mathrm{id}_{\emptyset_j})$$

generate respectively the cofibrations and the acyclic cofibrations of $\prod \mathscr{C}^{G_i}$ (see e.g. [Hir03, 11.1.10]), where $\emptyset_j$ is the initial object of $\mathscr{C}^{G_i}$. Moreover their images by L are precisely the families $F\mathcal{I}$ and $F\mathcal{I}$ from the statement. Following [Hir03, 11.3.1] and [Ste10, D.21], we prove that

i) $\prod \mathrm{ev}_j$ takes relative $L\mathcal{I}$-cell complexes to cofibrations: Let λ be a non-zero ordinal and $X\colon \lambda \to \mathscr{C}_a^I$ a functor such that for all morphism $\beta \to \beta'$ in λ the map $X_\beta \to X_{\beta'}$ is a pushout of a map in $L\mathcal{I}$. We need to show that for every $j \in I$ the map

$$\mathrm{ev}_j X_0 \longrightarrow \mathrm{ev}_j \operatorname*{colim}_\lambda X = \operatorname*{colim}_\lambda \mathrm{ev}_j \circ X$$

is a cofibration in $\mathscr{C}^{G_i}$. Since ev_j commutes with colimits, each map $\mathrm{ev}_j X_\beta \to \mathrm{ev}_j X_{\beta'}$ is the pushout of a map in $\mathrm{ev}_j L\mathcal{I}$. Thus we need to show that every map in $\mathrm{ev}_j L\mathcal{I}$ is a cofibration of $\mathscr{C}^{G_j}$. By definition of $\mathcal{I}$, this is the same as showing that for all $i, j \in I$ every generating cofibration of $\mathcal{I}_i$ is sent by $\mathrm{ev}_j F_i$ to a cofibration of G_j-C. The composite functor $\mathrm{ev}_j F_i$ is by definition

$$\mathrm{ev}_j F_i = K_{ji} \otimes_{G_i} (-)\colon \mathscr{C}^{G_i} \longrightarrow \mathscr{C}^{G_j}$$

which sends generating cofibrations to cofibrations by hypothesis (2.1).

ii) $\prod \mathrm{ev}_j$ takes relative $L\mathcal{J}$-cell complexes to acyclic cofibrations: the argument is similar to the one above.

Moreover $\prod \mathrm{ev}_j$ preserves colimits. By [Hir03, 11.3.1] and [Ste10, D.21], the families $L\mathcal{I}$ and $L\mathcal{J}$ are respectively a class of generating cofibrations and acyclic cofibrations for a $sSet^G$-enriched model structure on $\mathscr{C}_a^I$ with the fibrations and weak equivalences of the statement. □

Remark 2.9. Recall the isomorphism $\mathscr{C}_a^I \cong \mathscr{C}^{G \ltimes I}$ of Lemma 1.8. The model structure on $\mathscr{C}_a^I$ does not correspond to the projective model structure on $\mathscr{C}^{G \ltimes I}$. However, every fibration (resp. weak equivalence) in $\mathscr{C}_a^I$ is in particular a fibration (resp. weak equivalence) in $\mathscr{C}^{G \ltimes I}$. This means that the cofibrations of $\mathscr{C}^{G \ltimes I}$ are also cofibrations in $\mathscr{C}_a^I$. In particular, a sufficient condition for an object of $\mathscr{C}_a^I$ to be cofibrant is to be cofibrant in the projective model structure of $\mathscr{C}^{G \ltimes I}$.

Proposition 2.10: *If $X \in \mathscr{C}_a^I$ is cofibrant, each vertex X_i is cofibrant in $\mathscr{C}^{G_i}$.*

Proof. An argument analogous to the one in the proof of Lemma 2.8 shows that the right adjoint R_i to the evaluation functor $\mathrm{ev}_i\colon \mathscr{C}_a^I \to \mathscr{C}^{G_i}$ has j-vertex

$$\mathrm{ev}_j R_i = \hom_{G_i}(K_{ji}^*, -)$$

where K_{ji}^* is the set K_{ji} with left G_i-action $g \cdot k := k \cdot g^{-1}$ and right G_j-action $k \cdot g := g^{-1} \cdot k$. Since the fibrations and the equivalences on $\mathscr{C}_a^I$ are point-wise, $R_i\colon \mathscr{C}^{G_i} \to \mathscr{C}_a^I$ is a right Quillen functor. Therefore ev_i is a left Quillen functor, and in particular it preserves cofibrant objects. □

Definition 2.11. Let $\mathscr{C}$ and $\mathscr{D}$ be G-model categories. A G-Quillen adjunction (resp. equivalence) is an enriched adjunction $\mathscr{C} \rightleftarrows \mathscr{D}$ such that the induced adjunction $\mathscr{C}^H \rightleftarrows \mathscr{D}^H$ is a Quillen adjunction (resp. equivalence) for every subgroup $H \leq G$.

Example 2.12. The Quillen equivalence $|-|\colon sSet \rightleftarrows Top\colon Sing$ (see [GJ09, I]) is a G-Quillen equivalence for any finite group G.

Corollary 2.13: *A G-Quillen equivalence $L\colon \mathscr{C} \rightleftarrows \mathscr{D}\colon R$ induces a Quillen equivalence*

$$L\colon \mathscr{C}_a^I \rightleftarrows \mathscr{D}_a^I\colon R.$$

Proof. The adjunction $L\colon \mathscr{C}_a^I \rightleftarrows \mathscr{D}_a^I\colon R$ is a Quillen adjunction since the right adjoint preserves fibrations and acyclic fibrations, as they are defined point-wise. Let $X \in \mathscr{C}_a^I$ be cofibrant and $Y \in \mathscr{D}_a^I$ fibrant. A map $X \to R(Y)$ is an equivalence if and only if its adjoint $L(X) \to Y$ is, since by Proposition 2.10 X is point-wise cofibrant. □

2.3 Cofibrant replacement of G-diagrams

When $\mathscr{C}$ is a cofibrantly generated simplicial model category and I is a small category a standard way to replace a diagram $X\colon I \to \mathscr{C}$ by a cofibrant diagram is by the construction of Example 1.20. Namely, one defines qX by $qX_i = \mathrm{hocolim}_{I/i}(u_i^* X)$ where $u_i\colon I/i \to I$ is the functor that forgets the map to i. Then qX is cofibrant in the projective model structure on $\mathscr{C}^I$ and the natural map $\rho_X\colon qX \to X$ is a weak equivalence if X has cofibrant values in $\mathscr{C}$. In this section we will generalize this to G-diagrams as follows:

Theorem 2.14: *If X is a G-diagram such that for all i in I the value X_i is cofibrant in $\mathscr{C}^{G_i}$, then the map $\rho_X\colon qX \to X$ is a cofibrant replacement of G-diagrams.*

The proof is technical and will occupy the rest of this section. We begin by fixing some notation. Let I be a small category with an action a of G. Write I^δ for the discrete category with the same objects as I but no non-identity morphisms. The inclusion $I^\delta \hookrightarrow I$ is equivariant and induces a restriction functor $r\colon \mathscr{C}_a^I \to \mathscr{C}_a^{I^\delta}$ with left adjoint r_*. We abbreviate $r(X)$ as X^δ. Note that the functor r preserves fibrations and weak equivalences and so it is a right Quillen functor. It follows that the left adjoint r_* is a left Quillen functor. We say that an I-indexed G-diagram X is *point-wise cofibrant* if for each object i in I the value X_i is cofibrant in $\mathscr{C}^{G_i}$.

Lemma 2.15: *i) If Y is an I^δ-indexed G-diagram which is point-wise cofibrant, then Y is cofibrant in $\mathscr{C}_a^{I^\delta}$.*

ii) In particular, if X is a point-wise cofibrant I-indexed G-diagram then $r_* X^\delta$ is cofibrant in $\mathscr{C}_a^I$.

Proof. To see that part i) holds, consider a square

$$\begin{array}{ccc} \emptyset & \longrightarrow & Z \\ \downarrow & & \sim\downarrow f \\ Y & \longrightarrow & W \end{array} \tag{5}$$

in $\mathscr{C}_a^{I^\delta}$, where the right hand vertical map is a trivial fibration and $\emptyset$ denotes the initial object. The map f being a trivial fibration means exactly that each component $f_i\colon Z_i \to W_i$ is a trivial fibration in $\mathscr{C}^{G_i}$. Choose a representative i of each G-orbit in obI. Each resulting square

$$\begin{array}{ccc} \emptyset & \longrightarrow & Z_i \\ \downarrow & \overset{\lambda_i}{\nearrow} & \sim\downarrow f_i \\ Y_i & \longrightarrow & W_i \end{array}$$

has a lift λ_i since Y_i is cofibrant and f_i is a trivial fibration in $\mathscr{C}^{G_i}$. For $g \in G$ define $\lambda_{gi} = g_{Z_i} \circ \lambda_i \circ g_{Y_i}^{-1}$. Then, if $gi = i$ the G_i-equivariance of the map λ_i says precisely that $\lambda_i = g_{Z_i} \circ \lambda_i \circ g_{Y_i}^{-1} = \lambda_{gi}$, so for all i and all $g \in G$ the map λ_{gi} is well-defined. It is now easy to see that the λ_{gi}'s assemble to a map of G-diagrams giving a lift in the square (5).

Part ii) follows immediately from part i) and the fact that r_* is a left Quillen functor and hence preserves cofibrancy of objects. □

The adjunction (r_*, r) induces a comonad $r_* r$ on $\mathscr{C}_a^I$ in the usual way. For a G-diagram X the value $(r_* r)X$ on i is

$$(r_* r)X_i = \coprod_{f\colon j \to i} X_j.$$

The counit $\varepsilon\colon (r_*r)X \to X$ maps the X_j-component in the coproduct indexed by $f\colon j \to i$ to X_i by the map $X(f)$. The comultiplication $c\colon (r_*r)X \to (r_*rr_*r)X$ has as i-component the map

$$\coprod_{f\colon j\to i} X_j \to \coprod_{f\colon j\to i} \left(\coprod_{f'\colon k\to j} X_k \right)$$

that maps the X_j-summand indexed by $f\colon j \to i$ by the identity to the X_j-summand indexed by id_j in the f-summand of the target.

Let X be a G-diagram indexed on I. The bar construction on the comonad r_*r gives a simplicial G-diagram $B(r_*r)X$ with $B_n(r_*r)X = (r_*r)^{n+1}X$ so that

$$B_n(r_*r)X_i = \coprod_{f_0\colon i_0\to i} \coprod_{f_1\colon i_1\to i_0} \cdots \coprod_{f_n\colon i_n\to i_{n-1}} X_{i_n} \cong \coprod_{i_n\to\cdots\to i_0\to i} X_{i_n}.$$

Note that the indexing G_i-set can be identified with $N_n(I/i)^{op}$. For

$$\sigma = i_n \xrightarrow{\alpha_n} \cdots \xrightarrow{\alpha_1} i_0 \xrightarrow{\alpha_0} i$$

in $N_n(I/i)^{op}$ the face map d_k for $k > 0$ composes the maps α_k and α_{k-1} and d_n maps X_{i_n} to the $X_{i_{n-1}}$ indexed by $d_n(\sigma) \in N_{n-1}(I/i)^{op}$ by the map $X(\alpha_n)$. The degeneracy map s_l inserts an identity in the l-spot. Note that

$$\operatorname*{colim}_I r_*rX = \operatorname*{colim}_{I^\delta} rX = \coprod_i X_i,$$

so that $\operatorname{colim}_I B_n(r_*r)X \cong \coprod_{\sigma\in N_n(I^{op})} X_{\sigma(n)}$ and $\operatorname{colim}_I B(r_*r)X$ is isomorphic to the usual simplicial replacement $\coprod_* X$ of Bousfield and Kan [BK72] with G-action induced by the G-structure on X.

Proposition 2.16: *Let X be an I-indexed G-diagram. Then there are natural isomorphisms in $\mathscr{C}^G$*

i) $N(-/I)^{op} \otimes_I^a X \cong |\coprod_* X|$

ii) $|\coprod_* X| \cong \operatorname{colim}_I qX$.

Proof. To see i we first decompose the tensor product as an iterated coend (cf. [Rie13, §6.6])

$$N(-/I)^{op} \otimes_I^a X = \int^i N(i/I)^{op} \otimes X_i \cong \int^i \left(\int^{[n]} \Delta^n \times N_n\,(i/I)^{op} \right) \otimes X_i.$$

Here and in the rest of the proof we leave it to the reader to check that this is compatible with the G-structures on the diagrams. Rearranging the parentheses and switching the order of the coends gives the isomorphic object

$$\int^{[n]} \int^i \Delta^n \otimes (N_n\,(i/I)^{op} \otimes X_i) \cong \int^{[n]} \Delta^n \otimes \left(\int^i \coprod_{i\to i_n\to\cdots\to i_0} X_i \right).$$

Now we analyze the latter $\int^i$-factor. It is a coend of the G-diagram $I^{op} \times I \to \mathscr{C}$ given by

$$(i,j) \mapsto \coprod_{i\to i_n\to\cdots\to i_0} X_j.$$

This is isomorphic to the diagram

$$(i,j) \mapsto \coprod_{i_n\to\cdots\to i_0} I(i,i_n) \otimes X_j$$

and we note that since coends commute with colimits there is an isomorphism

$$\int^i \coprod_{i_n \to \cdots \to i_0} I(i, i_n) \otimes X_i \cong \coprod_{i_n \to \cdots \to i_0} \int^i I(i, i_n) \otimes X_i.$$

Here we must be careful since the representable functor $I(-, i_n)$ is not itself a G-diagram, but the coproduct $\coprod_{\sigma \in N_n(I^{op})} I(-, \sigma(n))$ of representable functors is. Finally, we observe that $\int^i I(i, i_n) \otimes X_i \cong X_{i_n}$ so that

$$\int^{[n]} \Delta^n \otimes \left(\int^i \coprod_{i \to i_n \to \cdots \to i_0} X_i \right) \cong \int^{[n]} \Delta^n \otimes \left(\coprod_{i_n \to \cdots \to i_0} X_{i_n} \right) = |\coprod_* X|$$

To get the isomorphism in ii) we recall the isomorphism $\operatorname{colim}_I B(r_* r)X \cong \coprod_* X$. Since realization commutes with colimits, there are natural isomorphisms

$$|\coprod_* X| \cong |\operatorname*{colim}_I B(r_* r)X| \cong \operatorname*{colim}_I |B(r_* r)X|.$$

Evaluating at i gives

$$|B(r_* r)X|_i = \left| [n] \mapsto \coprod_{i_n \to \cdots \to i_0 \to i} X_{i_n} \right| \cong \operatorname*{hocolim}_{I/i} (u_i^* X)$$

where the last isomorphism is an instance of i) for the G_i-diagram $u_i^* X \colon I/i \to \mathscr{C}$. This gives an isomorphism

$$\operatorname*{colim}_I |B(r_* r)X| \cong \operatorname*{colim}_I qX.$$

□

Lemma 2.17: *If X is a point-wise cofibrant G-diagram, then the simplicial object $B(r_* r)X$ is Reedy cofibrant in $(\mathscr{C}_a^I)^{\Delta^{op}}$.*

Proof. Let $L = L_n B(r_* rX)$ be the n-th latching object of $B(r_* rX)$. The natural map

$$L_n B(r_* rX) \to B_n(r_* rX) = B$$

is at each i in I the inclusion of the summands indexed by the degenerate n-simplices in $N_n(I/i)^{op}$ into the coproduct over all n-simplices. Thus B decomposes as a coproduct $B = L \amalg N$ where the value of N at i is the coproduct indexed over all the *non*-degenerate simplices of the nerve. The decomposition is clearly compatible with the G-diagram structure on each factor. The diagram N is obtained by applying r_* to a pointwise cofibrant I^δ-indexed G-diagram and is therefore cofibrant. It follows that the map $L \to B$ is a cofibration. □

Corollary 2.18: *If X is a point-wise cofibrant G-diagram, then qX is cofibrant.*

Proof. We know from the proof of Proposition 2.16 that qX is the realization of the simplicial object $B(r_* r)X$ which is Reedy cofibrant by Lemma 2.17. Since realization takes Reedy cofibrant objects to cofibrant objects [GJ09, VII,3.6] it follows that qX is cofibrant. □

Example 2.19. Let $*_I$ be the I-indexed G-diagram with value the terminal object $*$ of $sSet$. Then $q(*_I)_i = \operatorname{hocolim}_{I/i}(*_{I/i}) \cong N(I/i)^{op}$, so that $q(*_I) \cong N(I/-)^{op}$ and similarly $q(*_{I^{op}}) \cong N(-/I)$. By Corollary 2.18 it follows that the diagrams $N(I/-)$ and $N(-/I)^{op}$ are cofibrant as G-diagrams since $*$ is cofibrant in $sSet^{G_i}$ for all i in I and taking opposite simplicial sets preserves cofibrations. Further, given categories I and J with G-actions a and b respectively and an equivariant functor $F \colon I \to J$, then since the left Kan extension F_* preserves cofibrancy the diagrams $N(F/-) \cong F_* N(I/-)$ and $N(-/F)^{op} \cong F_* N(-/I)^{op}$ are also cofibrant in $\mathscr{C}_b^J$.

Proof of Theorem 2.14. It only remains to see that the map ρ_X is a weak equivalence. For this we must show that for each i the map $\rho_{X_i}\colon \operatorname{hocolim}_{I/i} u_i^* X \to X_i$ is a weak equivalence in $\mathscr{C}^{G_i}$. The functor $\iota\colon * \to I/i$ sending the unique object to the terminal object is homotopy cofinal in the sense of Definition 2.23, so by Theorem 2.24 the map $X_i = \operatorname{hocolim}_* \iota^* u_i^* X \to \operatorname{hocolim}_{I/i} u_i^* X$ is a weak equivalence. Since it is also section to the map ρ_{X_i} it follows by the two out of three property that ρ_{X_i} is a weak equivalence as well. □

2.4 Homotopy invariance of map and tensor

In this section $\mathscr{C}$ is a G-model category and a is a G-action on a small category I.

Proposition 2.20: *Let $X \in \mathscr{C}_a^I$ be a G-diagram in $\mathscr{C}$. If X is fibrant, the functor*

$$map_I^a(-, X)\colon (sSet_a^I)^{op} \longrightarrow \mathscr{C}^G$$

preserves equivalences of cofibrant objects (in $sSet_a^I$). Dually, if X is point-wise cofibrant, the functor

$$(-) \otimes_I^a X\colon sSet_a^{I^{op}} \longrightarrow \mathscr{C}^G$$

preserves equivalences of cofibrant objects.

Proof. We prove the statement for map_I^a, the proof for $\otimes_I^a$ is similar. Let $K \to L$ be an equivalence of cofibrant diagrams in $sSet_a^I$. By Ken Brown's Lemma we can assume that $K \to L$ is a cofibration (cf. [Hir03, 7.7.1]). To show that the induced map is an equivalence, we need to solve the the lifting problem

$$\begin{array}{ccc} A & \longrightarrow & map_I^a(L, X) \\ {\scriptstyle\simeq}\downarrow & \nearrow & \downarrow \\ B & \longrightarrow & map_I^a(K, X) \end{array}$$

for every cofibration $A \to B$ in $\mathscr{C}^G$. Let $Map_{\mathscr{C}}(B, X)$ be the G-diagram in $sSet$ given by $i \mapsto Map_{\mathscr{C}}(B, X_i)$ and where the G-structure is given by the maps $Map_{\mathscr{C}}(g^{-1}, g_{X_i})\colon Map_{\mathscr{C}}(B, X_i) \to Map_{\mathscr{C}}(B, X_{gi})$. There is an equivariant adjunction isomorphism

$$\underline{\mathscr{C}}^G(B, map_I^a(L, X)) \cong \underline{sSet}_a^I(L, Map_{\mathscr{C}}(B, X)).$$

Therefore the lifting problem above is equivalent to the lifting problem in $sSet_a^I$

$$\begin{array}{ccc} K & \longrightarrow & Map_{\mathscr{C}}(B, X) \\ {\scriptstyle\simeq}\downarrow & \nearrow & \downarrow \\ L & \longrightarrow & Map_{\mathscr{C}}(A, X) \end{array}$$

This can be solved if $Map_{\mathscr{C}}(B, X) \to Map_{\mathscr{C}}(A, X)$ is a fibration in $sSet_a^I$, i.e., if for every object $i \in I$ the map $Map_{\mathscr{C}}(B, X_i) \to Map_{\mathscr{C}}(A, X_i)$ is a fibration of simplicial G_i-sets. By assumption X_i is fibrant in $\mathscr{C}^{G_i}$ and $A \to B$ is restricts to a cofibration in $\mathscr{C}^{G_i}$, so by axiom $SM7$ for the $sSet^{G_i}$-enriched model category $\mathscr{C}^{G_i}$ the map is a fibration. □

Proposition 2.21: *If K is a cofibrant diagram in $sSet_a^I$, the functor*

$$map_I^a(K, -)\colon \mathscr{C}_a^I \longrightarrow \mathscr{C}^G$$

preserves equivalences of fibrant objects. Dually if K is cofibrant in $sSet_a^{I^{op}}$, the functor

$$K \otimes_I^a (-)\colon \mathscr{C}_a^I \longrightarrow \mathscr{C}^G$$

preserves equivalences of point-wise cofibrant objects.

Proof. The proof is the same as for the non-equivariant case of [Hir03, 18.4], using the equivariant adjunctions as in the proof of 2.20. □

The following result generalizes Villarroel's result [VF04, 6.1]:

Corollary 2.22: *The functors* $\operatorname{holim}\colon \mathscr{C}_a^I \to \mathscr{C}^G$ *and* $\operatorname{hocolim}\colon \mathscr{C}_a^I \to \mathscr{C}^G$ *preserve equivalences between fibrant diagrams and point-wise cofibrant diagrams respectively.*

Proof. Recall that homotopy limits and homotopy colimits are defined by cotensoring with $N(I/-)$ and tensoring with $N(-/I)^{op}$, respectively. By Proposition 2.21 it is enough to show that $N(I/-)$ is cofibrant in $sSet_a^I$ and $N(-/I)$ is cofibrant in $sSet_a^{I^{op}}$. This was shown in Example 2.19. □

2.5 Equivariant cofinality

Let I and J be categories with G-actions a and b respectively, $F\colon I \to J$ an equivariant functor, and $X\colon J \to \mathscr{C}$ a G-diagram.

We want to know when the canonical maps

$$\operatorname*{hocolim}_I F^*X \longrightarrow \operatorname*{hocolim}_J X \qquad \text{and} \qquad \operatorname*{holim}_J X \longrightarrow \operatorname*{holim}_I F^*X$$

are equivalences in $\mathscr{C}^G$. As in the non-equivariant setting, the categories F/j and j/F play a role in answering this question. For every object $j \in J$ they inherit a canonical action by the stabilizers group $G_j \leq G$ of j.

Definition 2.23. The functor $F\colon I \to J$ is left (resp. right) cofinal if for every $j \in J$ the nerve of the category F/j (resp. j/F) is weakly G_j-contractible.

Notice that for $H \leq G_i$, the H-fixed points of the nerve of F/j are isomorphic to the nerve of $(F/j)^H$. Therefore F is left cofinal if and only if the fixed categories $(F/j)^H$ are contractible for all $H \leq G_i$, and similarly for right cofinality.

The following is a generalization of [TW91, 1] and [VF04, 6.3].

Theorem 2.24: *Let* $\mathscr{C}$ *be a* G*-model category,* $F\colon I \to J$ *be an equivariant functor, and* $X \in \mathscr{C}_b^J$ *a* G*-diagram in* $\mathscr{C}$*. If* F *is left cofinal and* X *is fibrant, the canonical map*

$$\operatorname*{holim}_J X \longrightarrow \operatorname*{holim}_I F^*X$$

is an equivalence in $\mathscr{C}^G$*. Dually, if* F *is right cofinal and* X *is point-wise cofibrant, the map*

$$\operatorname*{hocolim}_I F^*X \longrightarrow \operatorname*{hocolim}_J X$$

is an equivalence in $\mathscr{C}^G$*.*

Proof. We prove the part of the statement about left cofinality. The map $\operatorname{holim}_J X \to \operatorname{holim}_I F^*X$ factors as

$$\begin{array}{ccc} map_J^b(NJ/(-), X) & \longrightarrow & map_I^a(NI/(-), F^*X) \\ & \searrow & \uparrow \cong \\ & & map_J^b(NF/(-), X) \end{array}$$

The vertical map is a cotensor version of the (F_*, F^*)-adjunction isomorphism. It is equivariant and it is showed to be an isomorphism in [Hir03, 19.6.6]. The diagonal map is induced by the projection map $NF/(-) \to NJ/(-)$ which is an equivalence in $sSet_b^J$, since for all $H \leq G$ and all object $j \in J^H$ both categories F/j^H and J/j^H are contractible (J/j^H has a final object). Moreover the G-diagrams $NJ/(-)$ and $NF/(-)$ are cofibrant in $sSet_a^J$, by Example 2.19. Therefore the induced map on homotopy limits is an equivalence by the homotopy invariance of map_J^b of Proposition 2.20. □

2.6 The Elmendorf theorem for G-diagrams

Let $\mathscr{C}$ be a cofibrantly generated model category with cellular fixed points, in the sense of [GM13]. Then the category $\mathscr{C}^G$ of G-object admits the fixed points model structure, where weak equivalences and fibrations are the equivariant maps whose H-fixed points are respectively weak equivalences and fibrations in $\mathscr{C}$, for every subgroup $H \leq G$. Let $\mathcal{O}_G$ be the orbit category of G, with quotient sets G/H as objects and equivariant maps as morphisms. Elmendorf's theorem (see [Ste10],[Elm83]) describes a Quillen equivalence

$$L\colon \mathscr{C}^{\mathcal{O}_G^{op}} \rightleftarrows \mathscr{C}^G\colon R$$

where the diagram category $\mathscr{C}^{\mathcal{O}_G^{op}}$ has the projective model structure. In this section we prove an analogous result, giving a Quillen equivalence between the category of G-diagrams in $\mathscr{C}$ and a category of diagrams with the projective model structure.

Let I be a small category with an action a of G. For convenience we will consider the category of G-diagrams in $\mathscr{C}$ as the category $\mathscr{C}^{G\rtimes_a I}$ of diagrams indexed over the Grothendieck construction of the action (see 1.8). The functor $a\colon G \to Cat$ induces a functor $\overline{a}\colon \mathcal{O}_G^{op} \to Cat$ that sends G/H to the category I^H of objects and morphisms of I fixed by the H-action. We denote its Grothendieck construction by $\mathcal{O}_G^{op} \rtimes_{\overline{a}} I$. The inclusion functor $G \to \mathcal{O}_G^{op}$ that sends the unique object to $G/1$ induces a functor $G \rtimes_a I \to \mathcal{O}_G^{op} \rtimes_{\overline{a}} I$. By restricting along this, we get a functor

$$L\colon \mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I} \longrightarrow \mathscr{C}^{G\rtimes_a I}$$

Recall from 2.4 that if the fixed point functors of $\mathscr{C}$ are cellular, the fixed points model structures on $\mathscr{C}^H$, for $H \leq G$, assemble into a G-model category.

Theorem 2.25: *Let $\mathscr{C}$ be a category such that the fixed points functors for the subgroups of G are cellular. The functor $L\colon \mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I} \to \mathscr{C}^{G\rtimes_a I}$ is the left adjoint of a Quillen equivalence*

$$L\colon \mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I} \rightleftarrows \mathscr{C}^{G\rtimes_a I}\colon R$$

where $\mathscr{C}^{G\rtimes_a I}$ has the model structure of 2.6 and $\mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I}$ has the projective model structure.

Proof. The right adjoint sends a G-diagram X in $\mathscr{C}^{G\rtimes_a I} \cong \mathscr{C}_a^I$ to the diagram $R(X)\colon \mathcal{O}_G^{op} \rtimes_{\overline{a}} I \to \mathscr{C}$ that sends an object $(G/H, i \in I^H)$ to

$$R(X)(G/H, i \in I^H) = X_i^H$$

In order to define $R(X)$ on morphisms, recall that the set of equivariant maps $G/K \to G/H$ is in natural bijection with $(G/H)^K$. A morphism in $\mathcal{O}_G^{op}$ from $(G/H, i)$ to $(G/K, j)$ is a pair $(z \in (G/H)^K, (\alpha\colon zi \to j) \in I^K)$, and it is sent to the composite

$$X_i^H \xrightarrow{z} X_{zi}^K \xrightarrow{\alpha_*^K} X_j^K$$

A morphisms $f\colon X \to Y$ in $\mathscr{C}_a^I$ is sent to the natural transformation with value $X_i^H \xrightarrow{f_i^H} Y_i^H$ at the object $(G/H, i \in I^H)$. It is straightforward to see that R is a right adjoint for L. The counit $LRX \to X$ is an isomorphism, and the unit at a diagram Z of $\mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I}$ is the natural transformation

$$\eta_Z\colon Z(G/H, i) \longrightarrow RL(Z)(G/H, i) = Z(G/1, i)^H$$

induced by the morphism $(H \in (G/H)^1, \mathrm{id}_i)\colon (G/H, i) \to (G/1, i)$ of $\mathcal{O}_G^{op} \rtimes_{\overline{a}} I$. By definition of the fixed points model structure and of the model structure on $\mathscr{C}^{G\rtimes_a I}$, the right adjoint R preserves and detects equivalences and fibrations. Thus the adjunction (L, R) is a Quillen pair.

Since R preserves and detects equivalences, (L, R) is a Quillen equivalence precisely if the unit $\eta_Z\colon Z \to RL(Z)$ is an equivalence for all cofibrant objects Z in $\mathscr{C}^{\mathcal{O}_G^{op}\rtimes_{\overline{a}} I}$. We prove this, following

[Ste10]. By cellularity of the fixed point functors RL preserves pushouts along generating cofibrations and directed colimits along point-wise cofibrations. Thus it is enough to show that η_Z is an isomorphism when Z is a generating cofibrant object, that is, an object of the form

$$Z = \hom_{\mathcal{O}_G^{op} \rtimes_{\overline{a}} I}((G/H, i), -) \otimes c$$

for fixed objects $(G/H, i)$ of $\mathcal{O}_G^{op} \rtimes_{\overline{a}} I$ and c of $\mathscr{C}$ cofibrant. For such a Z, the unit at an object $(G/K, j)$ is the top horizontal map of the commutative diagram

$$\begin{array}{ccc}
\hom_{\mathcal{O}_G^{op} \rtimes_{\overline{a}} I}((G/H, i), (G/K, j)) \otimes c & \xrightarrow{\eta} & (\hom_{\mathcal{O}_G^{op} \rtimes_{\overline{a}} I}((G/H, i), (G/e, j)) \otimes c)^K \\
\cong \downarrow & & \downarrow \cong \\
\{(z \in (G/H)^K, \alpha \colon (zi \to j) \in I^K)\} \otimes c & \longrightarrow & \big(\{(z \in G/H, \alpha \colon (zi \to j) \in I)\} \otimes c\big)^K \\
\| & & \| \\
\Lambda_{ij}^K \otimes c & \longrightarrow & (\Lambda_{ij} \otimes c)^K
\end{array}$$

where Λ_{ij} is the set of pairs $(z \in G/H, \alpha \in zi \to j)$ with K acting by left multiplication on G/H and by the category action on the map to j (notice that j belongs to I^K). The bottom horizontal map is an isomorphism by the cellularity conditions on the K-fixed points functor. □

For the G-model category of spaces, the Elmendorf theorem gives a description of the fixed points of the homotopy limit of a G-diagram as a space of natural transformations of diagrams.

Corollary 2.26: *For every G-diagram of spaces X in Top_a^I, there is a natural homeomorphism of spaces*

$$(\operatorname*{holim}_I X)^G \cong Map_{Top^{\mathcal{O}_G^{op} \rtimes_{\overline{a}} I}}\big(R(BI/(-)), R(X)\big)$$

where $R(X) \colon \mathcal{O}_G^{op} \rtimes_{\overline{a}} I \to Top$ has vertices $R(X)_{(G/H,i)} = X_i^H$.

Proof. The space $(\operatorname{holim}_I X)^G$ is by definition the mapping space from $BI/(-)$ to X in Top_a^I. As the counit of the adjunction of the Elmendorf theorem is a sequence of natural homeomorphisms

$$Map_{Top_a^I}\big(BI/(-), X\big) \cong Map_{Top_a^I}\big(LR(BI/(-)), X\big) \cong Map_{Top^{\mathcal{O}_G^{op} \rtimes_{\overline{a}} I}}\big(R(BI/(-)), R(X)\big)$$

□

3 Equivariant excision

3.1 Equivariant cubes and G-excision

If J is a finite G-set, the poset category of subsets $\mathcal{P}(J)$ has a canonical G-action, where a group element $g \in G$ sends a subset $U \subset J$ to the set

$$g \cdot U = \{g \cdot u \mid u \in U\}$$

Let $\mathscr{C}$ be a G-model category (cf. 2.1).

Definition 3.1. The category of J-cubes in $\mathscr{C}$ is the category of G-diagrams $\mathscr{C}_a^{\mathcal{P}(J)}$ for the action a on $\mathcal{P}(J)$ described above.

In order to define a homotopy invariant notion of (co)cartesian cubes, we need to make our homotopy (co)limits homotopy invariant. Given a cube $X \in \mathscr{C}_a^{\mathcal{P}(J)}$ let FX denote a fibrant J-cube together with an equivalence $X \xrightarrow{\sim} FX$. Similarly let $QX \xrightarrow{\sim} X$ denote an equivalence with QX point-wise cofibrant, that is, with QX_U cofibrant in $\mathscr{C}^{G_U}$ for every $U \in \mathcal{P}(J)$.

Remark 3.2. A replacement FX exists using the fibrant replacement in the model category $\mathscr{C}_a^{\mathcal{P}(J)}$. For the same reason one can find a replacement $QX \xrightarrow{\sim} X$ with QX cofibrant in $\mathscr{C}_a^{\mathcal{P}(J)}$, which is in particular point-wise cofibrant by 2.10. However, for a given cube one can often find a more explicit point-wise cofibrant replacement without having to find replacement cofibrant in $\mathscr{C}_a^{\mathcal{P}(J)}$ (see e.g. 3.4 and 3.5 below). For example, if a functorial cofibrant replacement Q in $\mathscr{C}$ lifts to a cofibrant replacement in $\mathscr{C}^H$ for every $H \leq G$, the diagram QX is point-wise cofibrant.

For any fixed object i of a category I with G-action, let $I\backslash i$ be the full subcategory of I on objects different from i, with the restricted G-action.

Definition 3.3. Let $\mathscr{C}$ be a G-model category. A J-cube $X \in \mathscr{C}_a^{\mathcal{P}(J)}$ is homotopy cocartesian if the canonical map

$$\operatorname*{hocolim}_{\mathcal{P}(J)\backslash J} i_J^* QX \longrightarrow QX_J \xrightarrow{\sim} X_J$$

is an equivalence in $\mathscr{C}^G$, where $i_J \colon \mathcal{P}(J)\backslash J \to \mathcal{P}(J)$ is the inclusion. Dually, $X \in \mathscr{C}_a^{\mathcal{P}(J)}$ is homotopy cartesian if the canonical map

$$X_\emptyset \xrightarrow{\sim} FX_\emptyset \longrightarrow \operatorname*{holim}_{\mathcal{P}(J)\backslash \emptyset} i_\emptyset^* FX$$

is an equivalence in $\mathscr{C}^G$, where $i_\emptyset \colon \mathcal{P}(J)\backslash\emptyset \to \mathcal{P}(J)$ is the inclusion.

Example 3.4. Let $c \in \mathscr{C}^G$ be a cofibrant object and J a finite G-set. Define a J_+-cube $S^J c$ by

$$(S^J c)_U = \begin{cases} c & , U = \emptyset \\ C^U c & , U \lneq J_+ \\ \Sigma^J c & , U = J_+ \end{cases}$$

Here $\Sigma^J c = \Sigma^{\tilde{J}_+} c$ is the suspension defined in 1.18, and $C^U c$ denotes the U-iterated cone

$$C^U c = \operatorname*{hocolim}_{\mathcal{P}(U)} \big(S \longmapsto \begin{cases} c & \text{if } S = \emptyset \\ * & \text{otherwise} \end{cases} \big) \simeq *$$

Since c is cofibrant, $S^J c$ is point-wise cofibrant. Moreover it is homotopy cocartesian. Indeed, its restriction to $\mathcal{P}(J_+)\backslash J_+$ is the cofibrant replacement q of Theorem 2.14 for the diagram $\sigma^J c \colon \mathcal{P}(J_+)\backslash J_+ \to \mathscr{C}$ with $(\sigma^J c)_\emptyset = c$ and the terminal object at the other vertices. Since homotopy colimits and colimits agree on cofibrant objects (by the homotopy invariance of $\otimes_I^a$), the canonical map from the homotopy colimit factors as

$$\operatorname*{hocolim}_{\mathcal{P}(J_+)\backslash J_+} S^J c = \operatorname*{hocolim}_{\mathcal{P}(J_+)\backslash J_+} q(\sigma^J c) \xrightarrow{\sim} \operatorname*{colim}_{\mathcal{P}(J_+)\backslash J_+} q(\sigma^J c) \cong \operatorname*{hocolim}_{\mathcal{P}(J_+)\backslash J_+} \sigma^J c = \Sigma^J c$$

Example 3.5. Suppose that $\mathscr{C}$ has a zero object, and denote the coproduct by $\bigvee$. Let $c \in \mathscr{C}^G$ be a cofibrant object and J a finite G-set. Define a J-cube $W^J c$ with vertices

$$(W^J c)_U = \begin{cases} \bigvee_J c & , U = \emptyset \\ c & , |U| = 1 \\ * & , |U| \geq 2 \end{cases}$$

with initial map $(W^J c)_\emptyset = \bigvee_J c \to c = (W^J c)_{\{j\}}$ the pinch map that collapses every wedge component different from j. This has a G-structure defined by the action on $\bigvee_J c$ on the initial vertex, and by the

action maps $g\colon (W^J c)_{\{j\}} = c \to c = (W^J c)_{\{gj\}}$. The cube Wc is homotopy cocartesian, that is, its homotopy colimit over $\mathcal{P}(J_+)\backslash J_+$ is equivalent in $\mathscr{C}^G$ to the zero object. To see this, we replace $W^J c$ by the equivalent cube

$$(\overline{W}c)_U = \begin{cases} \bigvee_J c & , U = \emptyset \\ c \bigvee_{J\backslash j} Cc & , U = \{j\} \\ \bigvee_J Cc & , |U| \geq 2 \end{cases}$$

where Cc is the one-fold cone $Cc = \mathrm{hocolim}(c \to *)$ and the non-identity maps of the diagram are all induced by cone inclusions $c \to Cc$. The G-structure is defined similarly as before, by permuting the wedge components. The cube $\overline{W}c$ is cofibrant, since the latching maps are all cofibrations (see A.1). As homotopy colimits preserve equivalences of point-wise cofibrant diagrams we get

$$\operatorname*{hocolim}_{\mathcal{P}(J_+)\backslash J_+} W^J c \xleftarrow{\simeq} \operatorname*{hocolim}_{\mathcal{P}(J_+)\backslash J_+} \overline{W}c \xrightarrow{\simeq} \operatorname*{colim}_{\mathcal{P}(J_+)\backslash J_+} \overline{W}c \cong \bigvee_J Cc$$

which is contractible since $\bigvee_J$ is a left Quillen functor.

We use the notion of homotopy cartesian and cocartesian G_+-cubes to express equivariant excision for functors between G-model categories $\mathscr{C}$ and $\mathscr{D}$. We shall consider functors for which we can express compatibility conditions with the model structures on $\mathscr{C}^H$ and $\mathscr{D}^H$ for every subgroup $H \leq G$, that is, functors $\Phi\colon \mathscr{C} \to \mathscr{D}^G$. A functor $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ induces a functor $\Phi_*\colon \mathscr{C}_a^I \to \mathscr{D}_a^I$ for any category with G-action I. The G-structure on $\Phi_*(X) = \Phi \circ X$ is defined by the maps

$$\Phi(X_i) \xrightarrow{g} \Phi(X_i) \xrightarrow{\Phi(g)} \Phi(X_{gi})$$

Since each map $\Phi(g)$ is G-equivariant $\Phi(g)g = g\Phi(g)$. For $I = *$ the trivial category this functor is the classical extension $\Phi_*\colon \mathscr{C}^G \to \mathscr{D}^G$. Similarly, the functor $\Phi\colon \mathscr{C} \to \mathscr{D}^H$ obtained by restricting the G-action to $H \leq G$, extends to a functor $\Phi_*\colon \mathscr{C}^H \to \mathscr{D}^H$

Definition 3.6. We call $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ a homotopy functor if for every subgroup $H \leq G$ the extended functor $\Phi_*\colon \mathscr{C}^H \to \mathscr{D}^H$ preserves equivalences of cofibrant objects. In particular the induced functor $\Phi_*\colon \mathscr{C}_a^I \to \mathscr{D}_a^I$ preserves equivalences of point-wise cofibrant G-diagrams.

Remark 3.7. The following are all examples of functors $\mathscr{C}^G \to \mathscr{D}^G$ that are extensions of homotopy functors $\mathscr{C} \to \mathscr{D}^G$.

- The identity functor $\mathscr{C}^G \to \mathscr{C}^G$,
- For a fixed pointed G-space K, the functors $K \wedge (-), Map_*(K, -)\colon Top_*^G \to Top_*^G$,
- Similarly for a fixed G-spectrum E the functor $E \wedge (-)\colon Top_*^G \to (\mathrm{Sp}^O)^G$.

An example of a functor $\mathscr{C}^G \to \mathscr{D}^G$ that is, not an extension of a functor $\mathscr{C} \to \mathscr{D}^G$ is the functor $(-)/G\colon Top^G \to Top^G$ that sends a G-space to its orbit space with trivial G-action.

Definition 3.8. Let $\mathscr{C}$ and $\mathscr{D}$ be pointed G-model categories. A homotopy functor $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ is called G-excisive if the induced functor $\Phi_*\colon \mathscr{C}_a^{\mathcal{P}(G_+)} \to \mathscr{D}_a^{\mathcal{P}(G_+)}$ sends homotopy cocartesian G_+-cubes to homotopy cartesian G_+-cubes. It is called G-linear if it is G-excisive and $\Phi(*)$ is equivalent to the zero object in $\mathscr{D}^G$.

Example 3.9. The following are examples of G-linear homotopy functors, as we will see later in the paper.

- Let M be an abelian group with additive G-action. Consider the homotopy functor $M(-)\colon sSet_* \to sSet_*^G$ that sends a simplicial set Z to

$$M(Z)_n = \bigoplus_{z \in Z_n} Mz/M*$$

where G acts diagonally on the direct summands. We show in 3.25 that this functor is G-linear, and explain how this is related to the equivariant Eilenberg-MacLane spectrum HM being a fibrant orthogonal G-spectrum.

- For a fixed G-spectrum $E \in (\mathrm{Sp}^O)^G$, the homotopy functor $E \wedge (-)\colon Top_* \to (\mathrm{Sp}^O)^G$ is G-linear (see 3.28).
- The inclusion as spectra with trivial G-action $\mathrm{Sp}^O \to (\mathrm{Sp}^O)^G$ (which extends to the identity on G-spectra) is G-linear (see 3.27).

Proposition 3.10: *A homotopy functor $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ is G-excisive if and only if the induced functor $\Phi_*\colon \mathscr{C}_a^{\mathcal{P}(G/H_+)} \to \mathscr{D}_a^{\mathcal{P}(G/H_+)}$ sends homotopy cocartesian G/H_+-cubes to homotopy cartesian G/H_+-cubes, for every subgroup $H \leq G$.*

Remark 3.11. Setting $H = G$ in 3.10 we see that $\Phi_*\colon \mathscr{C}_a^{\mathcal{P}(1_+)} \to \mathscr{D}_a^{\mathcal{P}(1_+)}$ sends cocartesian squares in $\mathscr{C}^G$ to cartesian squares in $\mathscr{D}^G$. That is, if Φ is G-excisive the induced functor $\Phi\colon \mathscr{C}^G \to \mathscr{D}^G$ is excisive in the classical sense.

Proof of 3.10. One of the implications of the statement is trivial. For the other one, there are two adjoint equivariant functors

$$p\colon \mathcal{P}(G_+) \rightleftarrows \mathcal{P}(G/H_+)\colon j$$

allowing us to compare homotopy limits and colimits of G_+- and G/H_+-cubes. They are both induced by the projection map $p\colon G_+ \to G/H_+$. The functor p sends $S \subset G_+$ to its image $p(S) = S \cdot H \subset G/H_+$, and j sends a subset $U \subset G/H_+$ to its preimage

$$j(U) = p^{-1}(U) = \bigcup_{u \in U} u \subset G_+$$

Notice that j is a section for p. The restriction $p_\emptyset\colon \mathcal{P}(G_+)\backslash\emptyset \to \mathcal{P}(G/H_+)\backslash\emptyset$ is left G-cofinal. Indeed, for any object $U \in \mathcal{P}(G/H_+)^K$ the subset $j(U) \subset G_+$ is final in $(p_\emptyset/U)^K$. Similarly, the restriction

$$j_{G/H_+}\colon \mathcal{P}(G/H_+)\backslash G/H_+ \longrightarrow \mathcal{P}(G_+)\backslash G_+$$

is right G-cofinal, the functor p providing initial objects for the categories $(S/j_{G/H_+})^K$.

Let $X\colon \mathcal{P}(G/H_+) \to \mathscr{C}$ be a homotopy cocartesian G/H_+-cube. The G_+-cube p^*X is also homotopy cocartesian, as there is a commutative diagram

$$\begin{array}{ccc}
\operatorname*{hocolim}_{\mathcal{P}(G_+)\backslash G_+} i_{G_+}^*(p^*QX) & \longrightarrow & (p^*QX)_{G_+} = QX_{G/H_+} \\
{\scriptstyle j_*}\uparrow\simeq & & \uparrow\simeq \\
\operatorname*{hocolim}_{\mathcal{P}(G/H_+)\backslash G/H_+} j_{G/H_+}^*(i_{G_+}^* p^* QX) & = & \operatorname*{hocolim}_{\mathcal{P}(G/H_+)\backslash G/H_+} i_{G/H_+}^* QX
\end{array}$$

The left vertical map is an equivalence by G-cofinality 2.24, and the right one since X is homotopy cocartesian. Notice that p^*QX is point-wise cofibrant as $G_S \subset G_{p(S)}$ and the forgetful functor $\mathscr{C}^{G_{p(S)}} \to \mathscr{C}^{G_S}$ preserves cofibrations. Therefore p^*QX is a point-wise cofibrant replacement of p^*X.

By G-excision, the G_+-cube $\Phi_* p^* X = p^* \Phi_* X$ is homotopy cartesian. There is a dual commutative diagram

$$\begin{array}{ccc}
\operatorname*{holim}_{\mathcal{P}(G/H_+)\backslash\emptyset} i_\emptyset^* F\Phi_* X & \longleftarrow & (F\Phi_* X)_\emptyset = (p^* F\Phi_* X)_\emptyset \\
{\scriptstyle p^*}\downarrow\simeq & & \downarrow\simeq \\
\operatorname*{holim}_{\mathcal{P}(G_+)\backslash\emptyset} p_\emptyset^*(i_\emptyset^* F\Phi_* X) & = & \operatorname*{holim}_{\mathcal{P}(G_+)\backslash\emptyset} i_\emptyset^* p^* F\Phi_* X
\end{array}$$

where the left vertical map is an equivalence by G-cofinality 2.24, and the right one since $p^*\Phi_* X$ is homotopy cartesian. This shows that $\Phi_* X$ is homotopy cartesian. □

Proposition 3.12: *Let $\mathscr{C}$ and $\mathscr{D}$ be pointed G-model categories, and $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ be a G-linear homotopy functor. For any finite G-set J and any cofibrant G-object $c \in \mathscr{C}^G$ the canonical map*

$$\Phi(\bigvee_J c) \longrightarrow \prod_J F\Phi(c)$$

is an equivalence $\mathscr{D}^G$.

Proof. First assume that $J = 1_+$ with trivial G-action. The square Vc

$$\begin{array}{ccc} c \vee c & \xrightarrow{p_+} & c \\ {\scriptstyle p_1}\downarrow & & \downarrow \\ c & \longrightarrow & * \end{array}$$

in $\mathscr{C}^G$ is homotopy cocartesian (cf. 3.5). By 3.11 its image $\Phi(Vc)$ is homotopy cartesian, that is, the map

$$\Phi(c \vee c) \xrightarrow{\simeq} F\Phi(c \vee c) \to \operatorname*{holim}_{\mathcal{P}(1_+)\setminus\emptyset} F\Phi(Vc) \simeq F\Phi(c) \times F\Phi(c)$$

is an equivalence in $\mathscr{D}^G$, with diagonal action on the target. By induction, the map of the statement is an equivalence for every J with trivial G-action. Given a finite G-set J, decompose it as disjoint union of transitive G-sets $J = \coprod_{z \in G\backslash J} z$. The map of the statement decomposes as

$$\Phi(\bigvee_J c) = \Phi(\bigvee_{z\in G\backslash J}\bigvee_z c) \xrightarrow{\simeq} \prod_{z\in G\backslash J} F\Phi(\bigvee_z c) \longrightarrow \prod_{z\in G\backslash J}\prod_z F\Phi(c) = \prod_J F\Phi(c)$$

with the first map an equivalence as the action on $G\backslash J$ is trivial. Therefore it is enough to show that the map is an equivalence for $J = G/H$ a transitive G-set.

Consider the G/H_+-cube Wc with vertices

$$(Wc)_U = \begin{cases} \bigvee_{G/H} c & , U = \emptyset \\ c & , U = \{j \neq +\} \\ * & , \text{otherwise} \end{cases}$$

It is homotopy cocartesian by an argument completely similar to 3.5. By 3.10 the cube $\Phi(Wc)$ is homotopy cartesian, that is, the canonical map

$$\Phi(\bigvee_{G/H} c) \to \operatorname*{holim}_{\mathcal{P}(G/H_+)\setminus\emptyset} F\Phi(Wc) \simeq \prod_{G/H} F\Phi(c)$$

is an equivalence in $\mathscr{D}^G$. □

Remark 3.13. In this equivariant setting G_+-cubes play the role that squares play in the classical theory. The equivariant analogue of n-cubes should be the category $\mathscr{C}_A^{\mathcal{P}(nG_+)}$, where nG is the disjoint union of n-copies of G with diagonal action. Following [Goo92], the behavior of Φ on nG_+-cubes X should be related to higher order G-excision.

3.2 The generalized Wirthmüller isomorphism

Let $\mathscr{C}$ be a bicomplete category, and G a finite group. We recall from §2.1 that a finite set K with commuting left H'-action and right H-action induces an adjunction

$$K \otimes_H (-)\colon \mathscr{C}^H \rightleftarrows \mathscr{C}^{H'}\colon \hom_{H'}(K,-)$$

Let K^* be the set K with left H-action and right H'-action defined by $h\cdot k\cdot h' = (h')^{-1}\cdot k\cdot h^{-1}$. If $\mathscr{C}$ is pointed (has an object that is, both initial and terminal) and if the actions on K are free, a functor $\Phi\colon \mathscr{C}\to\mathscr{D}^G$ induces a natural transformation

$$\eta\colon \Phi(K\otimes_H(-)) \longrightarrow \hom_H(K^*,\Phi(-))$$

of functors $\mathscr{C}^H\to\mathscr{D}^{H'}$. The map η_c is the image by the map

$$\begin{aligned}&\mathscr{C}^H((K^*\times_{H'}K)\otimes_H c, c)\xrightarrow{\Phi}\mathscr{D}^{H'}(\Phi((K^*\times_{H'}K)\otimes_H c),\Phi(c))\to\\ &\mathscr{D}^{H'}(K^*\otimes_{H'}\Phi(K\otimes_H c),\Phi(c))\xrightarrow{\cong}\mathscr{D}^{H'}(\Phi(K\otimes_H c),\hom_H(K^*,\Phi(c)))\end{aligned}$$

of the map $\coprod_{K^*\times_{H'}K_+} c\to c$ defined by $h\colon c\to c$ on a (k,k')-component with $k'h=k$, and by the trivial map $c\to *\to c$ otherwise. Notice that since the H-action is free there is at most one h for which $k'h=k$.

Example 3.14. Suppose that $K=G=H'$ with left G-multiplication and right H-multiplication. Sending an element to its inverse defines a H-G-equivariant isomorphism between G^* and G with left H-multiplication and right G-multiplication. We saw in 2.2 that the forgetful functor $\mathscr{C}^G\to\mathscr{C}^H$ is right adjoint to $G\otimes_H(-)$ and left adjoint to $\hom_H(G^*,-)$. The map η for the identity functor is the standard map

$$G\otimes_H(-)\longrightarrow\hom_H(G^*,-)$$

which in the case of spectra is the classical Wirthmüller isomorphism map. In 3.27 we apply 3.15 below to recover the Wirthmüller isomorphism theorem for G-spectra.

Theorem 3.15: *Let $\mathscr{C}$ and $\mathscr{D}$ be pointed G-model categories, and $\Phi\colon\mathscr{C}\to\mathscr{D}^G$ a G-linear homotopy functor. The composite*

$$\Phi(K\otimes_H c)\xrightarrow{\eta}\hom_H(K^*,\Phi(c))\longrightarrow\hom_H(K^*,F\Phi(c))$$

is an equivalence in $\mathscr{C}^{H'}$, where $\Phi(c)\xrightarrow{\simeq}F\Phi(c)$ is a fibrant replacement of $\Phi(c)$ in $\mathscr{D}^H$.

In particular if $\hom_H(K^*,-)$ preserves equivalences between all objects, the map $\eta\colon\Phi(K\otimes_H c)\to\hom_H(K^*,\Phi(c))$ is an equivalence for any $c\in\mathscr{C}^H$.

Proof. We define the map of the statement as a canonical map into a homotopy limit, and we use linearity of Φ to conclude that the map is an equivalence.

Choose a section $s_G\colon G/H\to G$ and a H'-H-equivariant map $\phi\colon K\to G$, which exists since the two actions on K are free (choose sections s and s' for the projections to K/H and $H'\backslash K$, and define $\phi(k)=h'h$ for $h\in H$ and $h'\in H'$ unique such that $k=s(kH)h=h's'(H'k)$). These choices give a commutative diagram (of sets)

$$\begin{array}{ccc} K & \xrightarrow{\phi} & G \\ {\scriptstyle s_K}\big(\big\downarrow{\scriptstyle \pi_K} & & {\scriptstyle \pi_G}\big\downarrow\big){\scriptstyle s_G} \\ K/H & \xrightarrow[\overline{\phi}]{} & G/H \end{array}$$

where $s_K(kH) = k \cdot (\phi(k)^{-1} \cdot s_G \pi_G \phi(k))$ is a section for π_K satisfying $\phi s_K = s_G \overline{\phi}$. This gives a map $\gamma\colon H' \times K/H \to H$ defined by

$$\gamma(h', z) = s_G(h'\overline{\phi}(z))^{-1} \cdot h' \cdot s_G \overline{\phi}(z)$$

inducing two functors $\bigvee_{K/H}(-)\colon \mathscr{C}^H \to \mathscr{C}^{H'}$ and $\prod_{K/H}(-)\colon \mathscr{D}^H \to \mathscr{D}^{H'}$ that send objects c and d respectively to the coproduct $\bigvee_{K/H} c$ and product $\prod_{K/H} d$ with respective H'-action

$$h'(z, x) = (h'z, \gamma(h', z) \cdot x) \qquad \text{and} \qquad (h' \cdot \underline{y})_z = \gamma(h', z) \cdot y_{(h')^{-1}z}$$

There is a commutative diagram of natural transformations

$$\begin{array}{ccccc} \Phi(\bigvee_{K/H} c) & \longrightarrow & \bigvee_{K/H} \Phi(c) & \longrightarrow & \prod_{K/H} \Phi(c) \\ \downarrow{\scriptstyle \Phi(s_K \otimes \mathrm{id}_c)}\,{\scriptstyle \cong} & & \downarrow{\scriptstyle s_K \otimes \mathrm{id}_{\Phi(c)}}\,{\scriptstyle \cong} & & \downarrow{\scriptstyle \cong}\,{\scriptstyle (-)\circ s_K} \\ \eta\colon \Phi(K \otimes_H c) & \longrightarrow & K \otimes_H \Phi(c) & \longrightarrow & \hom_H(K, \Phi(c)) \end{array}$$

where the top right horizontal map is the canonical map from the coproduct to the product. The map $s_K \otimes \mathrm{id}$ has inverse $(k, x) \mapsto (\pi_K k, (s_G \pi_G \phi(k))^{-1} \phi(k) \cdot x)$, and similarly for the right vertical map $(-) \circ s_K$. It is then enough to study the top composition $\Phi(\bigvee_{K/H} c) \to \prod_{K/H} \Phi(c)$.

Consider the K/H_+-cube $Wc\colon \mathcal{P}(K/H_+) \to \mathscr{C}$ defined by

$$(Wc)_S = \begin{cases} \bigvee_{K/H} c & , S = \emptyset \\ c & , |S| = 1, S \neq \{+\} \\ * & , |S| \geq 2 \text{ or } S = \{+\} \end{cases}$$

with initial map $\bigvee_{K/H} c \to c = (Wc)_z$ the pinch map that collapses all the wedge components non-indexed by z. The structure maps $c = (Wc)_z \to (Wc)_{h'z} = c$ are defined by action by $\gamma(h', z) \in H$.

The cube Wc is homotopy cocartesian. Indeed, if $Q_H c \xrightarrow{\simeq} c$ is a cofibrant replacement of c in $\mathscr{C}^H$, the cube $WQ_H c$ is point-wise cofibrant with homotopy colimit over $\mathcal{P}(K/H_+)\backslash K/H_+$ contractible (see 3.5). Let $\Phi(Wc) \xrightarrow{\simeq} F\Phi(Wc)$ be a fibrant replacement of $\Phi(Wc)$. By linearity of Φ, the canonical map

$$\Phi(\bigvee_{K/H} c) \xrightarrow{\simeq} \operatorname*{holim}_{\mathcal{P}(K/H_+)\backslash \emptyset} F\Phi(Wc) \xrightarrow{\simeq} \prod_{K/H} F\Phi(c)$$

is an equivalence in $\mathscr{D}^{H'}$. This map fits into a commutative diagram

$$\begin{array}{ccc} \Phi(\bigvee_{K/H} c) & \longrightarrow & \prod_{K/H} \Phi(c) \\ & \searrow^{\simeq} & \downarrow \\ & & \prod_{K/H} F\Phi(c) \end{array}$$

where the right vertical map is an equivalence if $\prod_{K/H}(-)$ preserves all equivalences. □

Corollary 3.16: *If the trivial action inclusion functor $\mathscr{C} \to \mathscr{C}^G$ is G-linear, the left and right adjoints to the evaluation functor $\mathrm{ev}_i\colon \mathscr{C}^I_a \to \mathscr{C}^{G_i}$ are naturally equivalent for every $i \in I$.*

Proof. We saw in 2.8 that the left adjoint $F_i\colon \mathscr{C}^{G_i} \to \mathscr{C}^I_a$ has j-vertex

$$(F_i c)_j = K_{ji} \otimes_{G_i} c$$

for $K_{ji} = \hom_{G \ltimes I}(i, j)$. Similarly the right adjoint has j-vertex

$$(R_i c)_j = \hom_{G_i}(K^*_{ji}, c)$$

and 3.15 provides a natural equivalence from F_i to R_i. □

3.3 G-linearity and adjoint assembly maps

Let $\mathcal{C}$ and $\mathcal{D}$ be G-model categories, and $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ a $sSet$-enriched reduced homotopy functor. Its extension $\Phi\colon \mathscr{C}^G \to \mathscr{D}^G$ is then enriched over G-$sSet$, and for any simplicial G-set K there is an assembly map

$$K \otimes \Phi(c) \longrightarrow \Phi(K \otimes c)$$

in $\mathscr{D}^G$. It is adjoint to the map of simplicial G-sets

$$K \longrightarrow Map_{\mathscr{C}}(c, K \otimes c) \xrightarrow{\Phi} Map_{\mathscr{C}}(\Phi(c), \Phi(K \otimes c))$$

where the first map is adjoint to the identity on $K \otimes c$. When $K = N(\mathcal{P}(J_+)\backslash\emptyset)$ this induces a map

$$\alpha\colon \Phi(c) \longrightarrow \Omega^J\Phi(\Sigma^J c)$$

called the adjoint assembly map.

Remark 3.17. Given a cofibrant G-object c in $\mathscr{C}^G$ and a finite G-set J, recall the cofibrant J_+-cube

$$(S^J c)_U = \begin{cases} c & , U = \emptyset \\ C^U c & , U \lneq J_+ \\ \Sigma^J c & , U = J_+ \end{cases}$$

from 3.4. This induces a map

$$\Phi(c) \xrightarrow{\simeq} F\Phi(c) \to \operatorname*{holim}_{\mathcal{P}(J_+)\backslash J_+} F\Phi(S^J c) \xleftarrow{\simeq} \Omega^J F\Phi(\Sigma^J c)$$

where the last equivalence is induced by the equivalence of fibrant $\mathcal{P}(J_+)\backslash J_+$-diagrams

$$\sigma^J\big(F\Phi(\Sigma^J c)\big) \xrightarrow{\simeq} F\Phi(S^J c)$$

where $\sigma^J d$ is the diagram from 1.18 associated to an object d of $\mathscr{D}^G$, with vertices $(\sigma^J d)_{J_+} = d$ and $(\sigma^J d)_U = *$ for $U \neq J$. The adjoint assembly map above fits into a diagram

$$\begin{array}{ccccc} \Phi(c) & \xrightarrow{\simeq} & F\Phi(c) & \longrightarrow & \operatorname*{holim}_{\mathcal{P}(J_+)\backslash J_+} F\Phi(S^J c) \\ & \searrow^{\alpha} & & & \uparrow{\scriptstyle \simeq} \\ & & \Omega^J\Phi(\Sigma^J c) & \longrightarrow & \Omega^J F\Phi(\Sigma^J c) \end{array}$$

Proposition 3.18: *Let $\mathscr{C}$ and $\mathscr{D}$ be pointed G-model categories, and $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ be a $sSet$-enriched G-linear homotopy functor. For any finite G-set J and any cofibrant G-object $c \in \mathscr{C}^G$ the composite*

$$\Phi(c) \xrightarrow{\alpha} \Omega^J\Phi(\Sigma^J c) \longrightarrow \Omega^J F\Phi(\Sigma^J c)$$

is an equivalence in $\mathscr{D}^G$.

Proof. The decomposition of J as disjoint union of transitive G-sets $J_+ = \bigvee_{z \in G\backslash J} z_+$ describes the map of the statement as an iterated construction

$$\Phi(c) \to \Omega^{z_1} F\Phi(\Sigma^{z_1} c) \to \cdots \to \Omega^{z_1} \ldots \Omega^{z_n} F\Phi(\Sigma^{z_1} \ldots \Sigma^{z_m} c)$$

Since $\Sigma^z(-)$ preserves cocartesian cubes and Ω^z preserves fibrant objects, it is enough to show that the map $\Phi(c) \to \Omega^{G/H} F\Phi(\Sigma^{G/H} c)$ is an equivalence for every transitive G-set G/H.

By 3.10, Φ sends the homotopy cocartesian G/H_+-cube $S^{G/H}c$ of 3.17 to a homotopy cartesian G/H_+-cube. that is, the map

$$\Phi(c) \xrightarrow{\simeq} F\Phi(c) \to \operatorname*{holim}_{\mathcal{P}(G/H_+)\backslash\emptyset} F\Phi(S^{G/H} c) \xleftarrow{\simeq} \Omega^{G/H} F\Phi(\Sigma^{G/H} c)$$

is an equivalence in $\mathscr{D}^G$. □

We aim at proving a converse for 3.18. We remind the reader that a simplicial category $\mathscr{C}$ is locally finitely presentable if there is a set of objects Θ in $\mathscr{C}$ such that every object of $\mathscr{C}$ is isomorphic to a filtered colimit of objects in Θ, and for every $\theta \in \Theta$ the functor $\mathrm{map}(\theta, -)\colon \mathscr{C} \to sSet$ preserves filtered colimits (see [AR94],[Kel82]). For example the categories of simplicial sets and spectra (of simplicial sets) satisfy this condition. For a finite G-set J and a subgroup $H \leq G$ we denote $J|_H$ the set J with action restricted to H.

Theorem 3.19: *Let $\mathscr{C}$ and $\mathscr{D}$ be pointed G-model categories, and suppose that $\mathscr{D}$ is G-Quillen equivalent to a G-model category $\mathscr{D}'$ with $(\mathscr{D}')^H$ finitely locally presentable for every $H \leq G$. Let $\Phi\colon \mathscr{C} \to \mathscr{D}^G$ be a $sSet$-enriched reduced homotopy functor, and J a finite G-set. If the canonical map*

$$\Phi(c) \longrightarrow \Omega^{J|_H} F\Phi(\Sigma^{J|_H} c)$$

is an equivalence in $\mathscr{D}^H$ for every cofibrant object $c \in \mathscr{C}^H$ and every subgroup $H \leq G$, then the induced functor $\Phi_*\colon \mathscr{C}_a^{\mathcal{P}(J_+)} \to \mathscr{D}_a^{\mathcal{P}(J_+)}$ sends homotopy cocartesian J_+-cubes to homotopy cartesian J_+-cubes.

In particular if $\Phi(c) \xrightarrow{\simeq} \Omega^{\rho|_H} F\Phi(\Sigma^{J|_H} c)$ is an equivalence for every subgroup $H \leq G$ and every cofibrant H-object c, the functor Φ is G-linear.

The proof of this theorem is given at the end of the section.

Corollary 3.20: *Under the hypotheses of 3.19, suppose additionally that the composite functor*

$$\mathscr{D}^H \xrightarrow{\mathrm{hom}_H(G_+,-)} \mathscr{D}^G \xrightarrow{(-)|_H} \mathscr{D}^H$$

detects equivalences. Then the following are equivalent:

1. *Φ is G-linear,*
2. *For every cofibrant object $c \in \mathscr{C}^H$, the canonical map $\Phi(c) \to \Omega^{\rho|_H} F\Phi(\Sigma^{\rho|_H} c)$ is an equivalence in $\mathscr{D}^H$,*
3. *For every finite G-set J the functor $\Phi_*\colon \mathscr{C}_a^{\mathcal{P}(J_+)} \to \mathscr{D}_a^{\mathcal{P}(J_+)}$ sends homotopy cocartesian J_+-cubes to homotopy cartesian J_+-cubes.*

Proof. (1) $\Rightarrow$ (2) Let c be a cofibrant object of $\mathscr{C}^H$. The G-object $G \otimes_H c$ is cofibrant in $\mathscr{C}^G$ as $G \otimes_H (-)$ is a left Quillen functor (see 2.1). Consider the homotopy cocartesian G_+-cube $S(G \otimes_H c)$ of 3.4 with vertices

$$S^G(G \otimes_H c)_U = \begin{cases} G \otimes_H c & , U = \emptyset \\ C^U(G \otimes_H c) & , U \lneq G_+ \\ \Sigma^\rho(G \otimes_H c) & , U = G_+ \end{cases}$$

By linearity of Φ the canonical map

$$\Phi(G \otimes_H c) \to \operatorname*{holim}_{\mathcal{P}(G_+)\setminus\emptyset} F\Phi(S^G(G \otimes_H c)) \xleftarrow{\simeq} \Omega^\rho F\Phi(\Sigma^\rho(G \otimes_H c))$$

is an equivalence in $\mathscr{D}^G$. Restricting the action to H, this map fits into a commutative diagram

$$\begin{array}{ccc} \Phi(G \otimes_H c)|_H & \xrightarrow{\simeq} & \Omega^{\rho|_H} F\Phi(\Sigma^\rho(G \otimes_H c))|_H \cong \Omega^{\rho|_H} F\Phi(G \otimes_H (\Sigma^{\rho|_H} c))|_H \\ \downarrow \simeq & & \downarrow \simeq \\ \mathrm{hom}_H(G_+, F\Phi(c))|_H & \longrightarrow & \mathrm{hom}_H(G_+, \Omega^{\rho|_H} F\Phi(\Sigma^{\rho|_H} c))|_H \cong \Omega^{\rho|_H} \mathrm{hom}_H(G_+, F\Phi(\Sigma^{\rho|_H} c))|_H \end{array}$$

where the vertical maps are the equivalences of 3.15 and the top horizontal map is an equivalence by 3.18. Thus the bottom horizontal map is also an equivalence, and since $\mathrm{hom}_H(G_+, -)|_H$ detects equivalences $\Phi(c) \to \Omega^{\rho|_H} F\Phi(\Sigma^{\rho|_H} c)$ is an equivalence as well.
(2) $\Rightarrow$ (3) This is 3.19.
(3) $\Rightarrow$ (1) This is 3.18. ∎

Remark 3.21. Define the G-derivative (at the point) of a reduced homotopy functor $\Phi\colon \mathcal{C} \to \mathcal{D}^G$ to be the functor $D_*\Phi\colon \mathcal{C} \to \mathcal{D}^G$ defined by

$$D_*\Phi(c) = \operatorname{hocolim}\big(Q\Phi(c) \to Q\Omega^\rho F\Phi(\Sigma^\rho c) \to Q\Omega^{2\rho} F\Phi(\Sigma^{2\rho} c) \to \dots\big)$$

where $n\rho$ is the direct sum of n copies of the regular representation of G. As a direct consequence of 3.20 the functor $D_*\Phi$ is G-linear, and it is equipped with a universal natural transformation $\Phi \to D_*\Phi$.

Proof of 3.19. Let $X\colon \mathcal{P}(J_+) \to \mathscr{C}$ be a cofibrant J_+-cube. For a cofibrant object $c \in \mathscr{C}^H$ let us denote

$$\overline{\Omega}^J F\Phi(\Sigma^J c) := \operatorname*{holim}_{\mathcal{P}(J_+)\backslash\emptyset} F\Phi(S^J c) \tilde{\leftarrow} \Omega^J F\Phi(\Sigma^J c)$$

Let $K\colon \mathcal{P}(J_+) \times \mathcal{P}(J_+) \to \mathscr{C}$ be the G-diagram

$$K(U,T) = \operatorname*{hocolim}_{S \in \mathcal{P}(J_+)\backslash J_+} X_{(S\cap U)\cup T}$$

and define a J_+-cube $Y\colon \mathcal{P}(J_+) \to \mathscr{D}$ by

$$Y_T = \operatorname*{holim}_{\mathcal{P}(J_+)\backslash\emptyset} F\Phi(K(-,T))$$

For every $T \subset J_+$, there is a commutative diagram, natural in T

$$\begin{array}{ccc} \Phi(X_T) & \longrightarrow & \overline{\Omega}^J F\Phi(\Sigma^J X_T) \\ & {\scriptstyle\phi}\searrow \quad \nearrow{\scriptstyle\psi} & \\ & Y_T & \end{array}$$

Denoting $\Delta^U := N\mathcal{P}(U)\backslash\emptyset$, the first map of the factorization has U-component

$$\phi_U\colon \Phi(X_T) \longrightarrow \hom(\Delta^U, F\Phi(K(U,T)))$$

adjoint to the composite

$$\Delta^U \otimes \Phi(X_T) \to \Phi(\Delta^U \otimes X_T) \to \Phi(K(U,T)) \to F\Phi(K(U,T))$$

where the second map is induced by $\Delta^U \otimes X_T \to \Delta^U \otimes X_{T\cup U} \to K(U,T)$. The map ψ is the homotopy limit over U of the map of diagrams $F\Phi(K(U,T)) \to F\Phi((S^J X_T)_U)$ induced by the map $K(U,T) \to (S^J X_T)_U$ defined as follows. For $U \neq J_+$, it is the composite

$$\begin{array}{ccccc} K(U,T) & = & \operatorname*{hocolim}_{S\in\mathcal{P}(J_+)\backslash J_+} X_{(S\cap U)\cup T} & = & \operatorname*{hocolim}_{S\in\mathcal{P}(J_+)\backslash J_+} (-\cap U)^* X_{S\cup T} \\ & \searrow & & & \downarrow \\ & & C^U X_T & = & \operatorname*{hocolim}_{S\in\mathcal{P}(U)} X_{S\cup T} \end{array}$$

where the vertical map is the canonical map induced by the functor $(-\cap U)\colon \mathcal{P}(J_+)\backslash J_+ \to \mathcal{P}(U)$. For $U = J_+$, the map is

$$K(J_+,T) = \operatorname*{hocolim}_{S\in\mathcal{P}(J_+)\backslash J_+} X_{S\cup T} \to \Sigma^J X_T$$

induced on homotopy colimits by the map of J_+-cubes given by the identity on the empty set vertex, and that collapses the other vertices to the point.

Now suppose that X is homotopy cocartesian, and let us see that Y is homotopy cartesian. There is a natural equivalence $K(U,T) \simeq X_{U\cup T}$. Indeed, the maps $X_{(S\cap U)\cup T} \to X_{((S\cup\{t\})\cap U)\cup T}$ are the identity for all $t \in T$, and therefore $K(U,T) \simeq X_{U\cup T}$ as long as $T \neq \emptyset$, by the lemma 3.22 below. For $T = \emptyset$ and $U \neq J_+$

$$K(U,\emptyset) = \operatorname*{hocolim}_{S\in\mathcal{P}(J_+)\setminus J_+} X_{S\cap U} \simeq X_U$$

again by 3.22, as the maps $X_{S\cap U} \to X_{(S\cup\{v\})\cap U}$ are the identity for all $v \in J_+\setminus U$. Finally,

$$K(J_+,\emptyset) = \operatorname*{hocolim}_{S\in\mathcal{P}(J_+)\setminus J_+} X_S \simeq X_{J_+}$$

since X is assumed to be homotopy cocartesian. This shows that

$$Y_T \simeq \operatorname*{holim}_{U\in\mathcal{P}(J_+)\setminus\emptyset} F\Phi(X_{U\cup T})$$

For every fixed $U \neq \emptyset$, the cube $T \longmapsto F\Phi(X_{U\cup T})$ is homotopy cartesian by 3.22, as the maps $F\Phi(X_{U\cup T}) \to F\Phi(X_{U\cup T\cup\{u\}})$ are the identity for all $u \in U$. The cube Y is then the homotopy limit of cartesian cubes, which is again cartesian as homotopy limits commute with each other.

Iterating this construction and using that Σ^J and Ω^J preserve respectively homotopy cocartesian and cartesian J_+-cubes, one gets a factorization of each map in the colimit system

$$\Phi(X) \xrightarrow{\simeq} \overline{\Omega}^J F\Phi(\Sigma^J X) \xrightarrow{\simeq} \overline{\Omega}^{2J} F\Phi(\Sigma^{2J} X) \xrightarrow{\simeq} \dots$$

through a homotopy cartesian J_+-cube $Y^{(n)}$. By (classical) cofinality for diagrams in $\mathscr{D}^G$ the homotopy colimit of the sequence above is equivalent to $\operatorname{hocolim}_n Y^{(n)}$. We prove in appendix A.3 that under our presentability assumptions sequential homotopy colimits preserve homotopy cartesian J_+-cubes. Therefore $\Phi(X) \simeq \operatorname{hocolim}_n Q\overline{\Omega}^{nJ} F\Phi(\Sigma^{nJ} X)$ is homotopy cartesian. □

Lemma 3.22: *Let J be a finite G-set, $X\colon \mathcal{P}(J) \to \mathscr{C}$ a J-cube and $I \subset J$ a non-empty G-invariant subset such that the maps $X_S \to X_{S\cup i}$ are isomorphisms for all $S \subset J$ and $i \in I$. If X is fibrant, it is homotopy cartesian. Similarly if X is point-wise cofibrant, it is homotopy cocartesian.*

Proof. Let $\mathcal{P}_I(J)$ be the subposet of non-empty subsets of J that contain I. By assumption, the restriction map

$$\operatorname*{holim}_{\mathcal{P}(J)\setminus\emptyset} X \longrightarrow \operatorname*{holim}_{\mathcal{P}_I(J)} X$$

is a G-equivariant homotopy equivalence. It fits into a commutative diagram

$$\begin{array}{ccc} X_\emptyset & \longrightarrow & \operatorname{holim}_{\mathcal{P}(J)\setminus\emptyset} X \\ \cong\downarrow & & \downarrow\simeq \\ X_I & \xrightarrow[\simeq]{} & \operatorname{holim}_{\mathcal{P}_I(J)} X \end{array}$$

The left vertical map is a G-map, which is an isomorphism by assumption. The bottom horizontal map is a G-equivalence since I is initial in $\mathcal{P}_I(J)$.

A completely analogous argument shows that X is homotopy cocartesian. □

3.4 G-linear functors on G-spaces

In [Blu06] Blumberg defines a notion of G-linearity for endofunctors of G-spaces, for a compact Lie group G. When G is finite, we show that his definition and ours agree up to a suspension factor.

Before starting, let us remark that when working with spaces we can drop all the point-wise fibrant and cofibrant replacements from the last sections, as homotopy limits and homotopy colimits of G-diagrams of spaces are always homotopy invariant. For homotopy limits, it is just because every G-space is fibrant. For homotopy colimits, there is a natural isomorphism

$$(\operatorname*{hocolim}_I X)^H \cong \operatorname*{hocolim}_{I^H}(\iota_H^* X)^H$$

for every G-diagram $X \in (Top_*)_a^I$ and $H \leq G$. Here $\iota_H \colon I^H \to I$ is the inclusion of the subcategory of I of objects and morphisms fixed by the H-action. Therefore homotopy invariance of homotopy colimits of G-diagrams follows from homotopy invariance of classical homotopy colimits of spaces, proved in [DI04].

Proposition 3.23: *An enriched reduced homotopy functor $\Phi \colon Top_* \to Top_*^G$ is G-linear if and only if the two following conditions hold*

a) The induced functor $\Phi_* \colon (Top_*^G)^{\mathcal{P}(1_+)} \to (Top_*^G)^{\mathcal{P}(1_+)}$ sends homotopy cocartesian squares of pointed G-spaces to homotopy cartesian ones,

b) For all finite G-set J the map

$$\Phi(\bigvee_J Z) \to \prod_J \Phi(Z)$$

is an equivalence of pointed G-spaces.

Remark 3.24. The two conditions of 3.23 are essentially the definition of G-linearity in the case of a finite group G in [Blu06].

Proof. If Φ is G-linear, it sends homotopy cocartesian squares to homotopy cartesian squares by 3.11, and the map $\Phi(\bigvee_J Z) \to \prod_J \Phi(Z)$ is an equivalence by 3.12.

Conversely, Blumberg proves in [Blu06] that the two conditions above imply that the adjoint assembly map $\Phi(Z) \to \Omega^V \Phi(Z \wedge S^V)$ is a G-equivalence for every G-representation V. By 3.19 this implies G-linearity of Φ. □

Example 3.25. Let M be a commutative topological monoid with additive G-action, and suppose that the fixed points monoids M^H are group-like for every subgroup H of G. The equivariant Dold-Thom construction $M(-) \colon Top_* \to Top_*^G$ sends a pointed space Z to the space $M(Z)$ of reduced configurations of points in Z with labels in M. Here G acts on the labels. After extending $M(-)$ to Top_*^G the group acts both on the labels and on the space. For a pointed G-simplicial set K, there is a G-homeomorphism $M(K) \cong M(|K|)$ where $M(K)$ is the construction of 3.9.

Using 3.23, we prove that $M(-) \colon Top_* \to Top_*^G$ is G-linear. Given a G-space Z, the fixed points of the map $M(Z) \to \Omega M(Z \wedge S^1)$ are described by the adjoint assembly map

$$M(Z)^H \longrightarrow \Omega M(Z \wedge S^1)^H \cong \Omega(M(Z))(S^1)^H \cong \Omega M(Z)^H(S^1)$$

for the topological group-like monoid $M(Z)^H$. This is an equivalence by standard arguments, see [May75, 7.6]. By 3.19 for the trivial G-set $J = \{1\}$, the functor $M(-)$ sends homotopy cocartesian squares of G-spaces to homotopy cartesian ones. Moreover the map $M(\bigvee_J Z) \to \prod_J M(Z)$ is an equivariant isomorphism. Thus $M(-) \colon Top_* \to Top_*^G$ is G-linear by 3.23.

Notice that by G-linearity the map $M(Z) \to \Omega^J M(Z \wedge S^J)$ is a G-equivalence for every finite G-set J. This shows that the associated Eilenberg-MacLane G-spectrum $HM_n = M(S^n)$ is fibrant in $(\mathrm{Sp}^O)^G$.

3.5 G-linear functors to G-spectra

Let us start by clarifying that when working with spectra, as for spaces, we can forget all about the point-wise cofibrant and fibrant replacements from the last sections, thanks to the following.

Lemma 3.26: *Let G be a finite group with an action $a\colon G \to Cat$ on the small category $a(*) = I$.*

- *The homotopy colimit functor* $\operatorname{hocolim}\colon (\mathrm{Sp}^O)^I_a \to (\mathrm{Sp}^O)^G$ *preserves equivalences between any two diagrams (not necessarily of cofibrant objects).*
- *If I has finite nerve, the homotopy limit functor* $\operatorname{holim}\colon (\mathrm{Sp}^O)^I_a \to (\mathrm{Sp}^O)^G$ *preserves equivalences between any diagram (not necessarily of fibrant objects).*

Proof. For any H-spectrum E there is a functorial cofibrant replacement $QE \to E$ where the map is a level equivalence. By 2.21 it is enough to show that homotopy colimits preserve level equivalences of maps of G-diagrams. Since homotopy colimits of spectra are defined level-wise, this follows from homotopy invariance of homotopy colimits for spaces (see §3.4).

For the statement about homotopy limits, take a G-diagram of spectra X. The positive equivariant homotopy groups of $\operatorname{holim}_I X$ are the homotopy groups of the G-space

$$\operatorname*{hocolim}_n \Omega^{n\rho}(\operatorname*{holim}_I X)(n\rho)$$

We used the notation $E(n\rho) = E_n \wedge_{O(n)} L(\mathbb{R}^{n|G|}, n\rho)^+$ for a G-spectrum E, where $L(\mathbb{R}^{n|G|}, n\rho)$ is the space of isomorphisms of vector spaces from $\mathbb{R}^{n|G|}$ to $n\rho$. There are natural equivalences

$$\begin{aligned}&\operatorname{hocolim}_n \Omega^{n\rho}(\operatorname{holim}_I X)(n\rho) \cong \operatorname{hocolim}_n \Omega^{n\rho} \operatorname{holim}_I X(n\rho) \cong\\ &\operatorname{hocolim}_n \operatorname{holim}_I \Omega^{n\rho} X(n\rho) \simeq \operatorname{holim}_I \operatorname{hocolim}_n \Omega^{n\rho} X(n\rho)\end{aligned}$$

where the last equivalence follows from A.3. Therefore an equivalence of G-diagrams of spectra $f\colon X \to Y$ induces an isomorphism in positive homotopy groups of the homotopy limit precisely when the map $\operatorname{holim}_I \operatorname{hocolim}_n \Omega^{n\rho} f^{(n\rho)}$ is an equivalence of G-spaces. Since f is an equivalence of G-diagrams of spectra, $\operatorname{hocolim}_n \Omega^{n\rho} f_i^{(n\rho)}$ is an equivalence of G_i-spaces for all $i \in I$. The map above is then the homotopy limit of an equivalence of G-diagrams of spaces, which is an equivalence by 2.20 since every G_i-space is fibrant. A similar argument shows that $\operatorname{holim}_I f$ is an equivalence in negative homotopy groups. □

Corollary 3.27: *Let J be a finite G-set and I a category with action A. Any homotopy cocartesian J_+-cube of spectra $X \in (\mathrm{Sp}^O)^{\mathcal{P}(J_+)}_a$ is homotopy cartesian, that is, the inclusion functor $\mathrm{Sp}^O \to (\mathrm{Sp}^O)^G$ is G-linear.*

In particular this implies the Wirthmüller isomorphism theorem, stating that for any subgroup $H \leq G$ and H-spectrum $E \in (\mathrm{Sp}^O)^H$ the canonical map

$$\eta\colon G \otimes_H E = G_+ \wedge_H E \longrightarrow \hom_H(G, E)$$

is an equivalence of G-spectra.

Proof. The category of G-spectra satisfies the presentability condition of 3.19. By the equivariant suspension theorem, the map $E \to \Omega^\rho(E \wedge S^\rho)$ is an equivalence for any H-spectrum E. By 3.19 this is equivalent to G-linearity of the functor $\mathrm{Sp}^O \to (\mathrm{Sp}^O)^G$. The map $\eta\colon G \otimes_H E \longrightarrow \hom_H(G, E)$ is an equivalence by 3.15, as $\hom_H(G, -)\colon (\mathrm{Sp}^O)^H \to (\mathrm{Sp}^O)^G$ preserves equivalences between all spectra. □

We end the section with a complete characterization of G-linear functors from the category $sSet^f_*$ of finite pointed simplicial sets to G-spectra.

Proposition 3.28: *Let $\Phi\colon sSet^f_* \to (\mathrm{Sp}^O)^G$ be a $sSet$-enriched reduced homotopy functor. The following conditions are equivalent*

1. *Φ is G-linear,*

2. $\Phi_*\colon ((sSet_*^f)^G)^{\mathcal{P}(1_+)} \to ((\mathrm{Sp}^O)^G)^{\mathcal{P}(1_+)}$ *sends homotopy cocartesian squares in* $(sSet_*^f)^G$ *to homotopy cartesian squares of* G*-spectra, and* $\Phi(\bigvee_J K) \to \prod_J \Phi(K)$ *is an equivalence for every finite pointed simplicial* G*-set* K *and finite* G*-set* J*,*

3. *For every* $K \in (sSet_*^f)^G$ *the assembly map*

$$\Phi(S^0) \wedge |K| \longrightarrow \Phi(K)$$

is an equivalence of G*-spectra.*

Proof. (1) $\Rightarrow$ (2) This is true in general, by 3.11 and 3.12.
(2) $\Rightarrow$ (3) can be proved by induction on the skeleton of K. The wedges into products condition gives the equivalence for the 0-skeleton, and the induction step follows from the condition on squares. We refer to [Dot13] for the details.
(3) $\Rightarrow$ (1) Since G-linearity is invariant under equivalences, we show that $E \wedge |-|$ is G-linear for any G-spectrum E. If $X\colon \mathcal{P}(G_+) \to sSet_*^f$ is homotopy cocartesian, the cube of spectra $E \wedge |X|$ is also homotopy cocartesian. Indeed, the geometric fixed points F^H of the canonical map from the homotopy colimit is

$$F^H(\operatorname*{hocolim}_{\mathcal{P}(G_+)\setminus G_+} E \wedge |X|) \cong F^H(E \wedge \operatorname*{hocolim}_{\mathcal{P}(G_+)\setminus G_+} |X|) \cong F^H(E) \wedge (\operatorname*{hocolim}_{\mathcal{P}(G_+)\setminus G_+} |X|)^H \xrightarrow{\simeq}$$
$$\xrightarrow{\simeq} F^H(E) \wedge |X_{G_+}|^H \cong F^H(E \wedge |X_{G_+}|)$$

This is an equivalence since X is homotopy cocartesian, and since smashing with a spectrum preserves homotopy equivalences of spaces. By 3.27 $E \wedge X$ is also homotopy cartesian. □

A Appendix

A.1 Finite categories and cofibrant G-diagrams

We give a criterion for determining when a G-diagram is cofibrant in the model structure of 2.6, when the over categories of the indexing category I have finite dimensional nerve. These are sometimes called directed Reedy categories. The criterion is in terms of latching maps, and it is completely analogous to the classical theory (see e.g. [Hir03][§15]).

Let $\mathscr{C}$ be a cocomplete category. We denote by $(I/i)'$ the over category I/i with the object $i = i$ removed. The latching diagram of a diagram $X\colon I \to \mathscr{C}$ is the diagram $L(X)\colon I \to \mathscr{C}$ with vertices

$$L(X)_i = \operatorname*{colim}_{(I/i)'}((I/i)' \longrightarrow I \xrightarrow{X} \mathscr{C})$$

and maps induced by the standard maps $(I/i)' \to (I/j)'$ associated to a map $i \to j$. This construction comes with a natural transformation $L(X) \to X$.

For a G-diagram $X \in \mathscr{C}_a^I$, the latching diagram $L(X)$ inherits a G-structure by the natural maps



and the map canonical map $L(X) \to X$ is a map of G-diagrams.

Proposition A.1: *Let $\mathscr{C}$ be a G-model category (see 2.1), and I a category with G-action such that the simplicial set NI/i is finite dimensional for every $i \in I$. Let X be an object of $\in \mathscr{C}_a^I$ such that the map $L(X)_i \to X_i$ is a cofibration in $\mathscr{C}^{G_i}$ for every $i \in I$. Then X is cofibrant in the model structure on $\mathscr{C}_a^I$ of 2.6.*

Proof. In order to show that X is cofibrant we need to define a lift for every diagram in $\mathscr{C}_a^I$

$$\begin{array}{ccc} & & Y \\ & \overset{l}{\nearrow} & \downarrow \sim \\ X & \longrightarrow & Z \end{array}$$

where the vertical map is an acyclic fibration. We build this lift by induction on a filtration of I defined by the degree function $\deg\colon ObI \to \mathbb{N}$ defined by

$$\deg(i) = \dim NI/i$$

It is easy to see that the degree function is equivariant (where $\mathbb{N}$ has trivial action), and that if $\alpha\colon i \to j$ is a non-identity morphism then $\deg(i) < \deg(j)$. Let $I_{\leq n}$ be the full subcategory of I with objects with degree less than or equal to n. Since the degree function is equivariant, the G-action of I restricts to $I_{\leq n}$, and the G-structure on X restricts to a G-structure on the restricted diagram $X_{\leq n}\colon I_{\leq n} \to I \xrightarrow{X} \mathscr{C}$. We build the lift inductively on the diagrams $X_{\leq n}$.

For the base step, choose a section $s\colon ObI_{\leq 0}/G \to ObI_{\leq 0}$. For every orbit $\gamma \in ObI_{\leq 0}/G$ one can choose a $G_{s(\gamma)}$-equivariant lift

$$\begin{array}{ccc} & & Y_{s(\gamma)} \\ & \overset{l_{s(\gamma)}}{\nearrow} & \downarrow \sim \\ X_{s(\gamma)} & \longrightarrow & Z_{s(\gamma)} \end{array}$$

since the map $\emptyset = L(X)_{s(\gamma)} \to X_{s(\gamma)}$ is a cofibration in $\mathscr{C}^{G_{s(\gamma)}}$ by assumption (the map $Y_{s(\gamma)} \to Z_{s(\gamma)}$ is an acyclic fibration of $\mathscr{C}^{G_{s(\gamma)}}$ by definition of the model structure on $\mathscr{C}_a^I$). Given any object $i \in I_{\leq 0}$ outside the image of s, define $l_i\colon X_i \to Y_i$ as the composite

$$X_i \xrightarrow{g^{-1}} X_{s([i])} \xrightarrow{l_{s[i]}} Y_{s([i])} \xrightarrow{g} Y_i$$

for a choice of $g \in G$ with $gs[i] = i$. Since the category $I_{\leq 0}$ is discrete (a G-set) by the properties of the degree function, these lifts define a map of diagrams $l^0\colon X_{\leq 0} \to Y_{\leq 0}$ lifting $X_{\leq 0} \to Z_{\leq 0}$. Moreover l respects the G-structure since the lifts $l_{s(\gamma)}$ are $G_{s(\gamma)}$-equivariant.

Now suppose we defined a lift $l^{n-1}\colon X_{\leq n-1} \to Y_{\leq n-1}$. Let I_n be the full subcategory of I on objects of degree n. Choose a section $s^n\colon ObI_n/G \to ObI_n$, and for every $\gamma \in ObI_n/G$ a lift in $\mathscr{C}^{G_{s^n(\gamma)}}$

$$\begin{array}{ccc} L(X)_{s^n(\gamma)} & \longrightarrow & Y_{s^n(\gamma)} \\ \downarrow & \overset{l_{s^n(\gamma)}}{\nearrow} & \downarrow \sim \\ X_{s^n(\gamma)} & \longrightarrow & Z_{s^n(\gamma)} \end{array}$$

The top horizontal map is the canonical map given by the universal property of the colimits defining $L(X)$. Again, the lifts exist because $L(X)_{s^n(\gamma)} \to X_{s^n(\gamma)}$ is a cofibration. For a general object i of I_n define

$$l_i\colon X_i \xrightarrow{g^{-1}} X_{s([i])} \xrightarrow{l_{s[i]}} Y_{s([i])} \xrightarrow{g} Y_i$$

Commutativity of the diagram above insures that the resulting map $l^n\colon X_{\leq n} \to Y_{\leq n}$commutes with the structure maps of $X_{\leq n}$ and $Y_{\leq n}$. Moreover l^n respects the G-structure by $G_{s^n(\gamma)}$-equivariancy of $l_{s(\gamma)}$. ∎

A.2 Sequential homotopy colimits and finite G-homotopy limits

Definition A.2 ([Kel82])**.** A simplicial category $\mathscr{C}$ is finitely locally presentable if there is a set of objects Θ satisfying

1. For every $c \in \Theta$ the mapping space functor
$$\mathrm{map}(c, -)\colon \mathscr{C} \longrightarrow sSet$$
preserves filtered colimits,
2. every object of $\mathscr{C}$ is isomorphic to a filtered colimit of objects in Θ.

We consider the poset category $\mathbb{N}$ of natural numbers as a category with trivial G-action.

Proposition A.3: *Let $\mathscr{C}$ be a G-model category, and suppose that the underlying simplicial categories $\mathscr{C}^H$ are finitely locally presentable for all $H \leq G$. Let J be a finite G-set and $X\colon \mathbb{N} \times \mathcal{P}(J_+) \to \mathscr{C}$ a G-diagram with the property that for every $n \in \mathbb{N}$ the J_+-cube X_n is homotopy cartesian. Then the J_+-cube* $\mathrm{hocolim}_{\mathbb{N}} QX_n$ *is also homotopy cartesian.*

Proof. We need to show that the top horizontal map in the commutative diagram

$$\begin{array}{ccc} \underset{\mathbb{N}}{\mathrm{hocolim}}\, QX_{n,\emptyset} & \longrightarrow & \underset{S\in\mathcal{P}(J_+)\setminus\emptyset}{\mathrm{holim}} F \underset{\mathbb{N}}{\mathrm{hocolim}}\, QX_{n,S} \\ \simeq\downarrow & & \downarrow\simeq \\ \underset{\mathbb{N}}{\mathrm{colim}}\, X_{n,\emptyset} & \longrightarrow & \underset{S\in\mathcal{P}(J_+)\setminus\emptyset}{\mathrm{holim}} F \underset{\mathbb{N}}{\mathrm{colim}}\, X_{n,S} \end{array}$$

is an equivalence in $\mathscr{C}^G$. The left vertical map is an equivalence since in the locally finitely presentable category $\mathscr{C}^G$ filtered colimits are homotopy colimits (see e.g. [Dug01, 7.3], or [BK72] for simplicial sets). Similarly, the right vertical map is the fibrant homotopy limit of an equivalence of G-diagrams, as each $\mathscr{C}^{G_S}$ is locally finitely presentable. The bottom map fits into a commutative diagram

$$\begin{array}{ccc} \underset{\mathbb{N}}{\mathrm{colim}}\, X_{n,\emptyset} & \longrightarrow & \underset{S\in\mathcal{P}(J_+)\setminus\emptyset}{\mathrm{holim}} F \underset{\mathbb{N}}{\mathrm{colim}}\, X_{n,S} \\ & \searrow^{\simeq} & \uparrow \\ & & \underset{\mathbb{N}}{\mathrm{colim}} \underset{S\in\mathcal{P}(J_+)\setminus\emptyset}{\mathrm{holim}} FX_{n,S} \end{array}$$

with the diagonal map an equivalence in $\mathscr{C}^G$ since X_n is homotopy cartesian and filtered colimits in $\mathscr{C}^G$ preserve equivalences. To show that the vertical map is an equivalence, we compute from the definition of homotopy limits. Denoting $K_S = N\mathcal{P}(S)\setminus\emptyset$ we have an isomorphisms in $\mathscr{C}^G$

$$\begin{aligned} &\underset{\mathbb{N}}{\mathrm{colim}} \underset{S\in\mathcal{P}(J_+)\setminus\emptyset}{\mathrm{holim}} FX_{n,S} = \underset{\mathbb{N}}{\mathrm{colim}}\, \lim \left(\prod_S map_{\mathscr{C}}(K_S, FX_{S,n}) \rightrightarrows \prod_{S\to T} map_{\mathscr{C}}(K_s, FX_{T,n}) \right) \cong \\ &\cong \lim \left(\prod_S \underset{\mathbb{N}}{\mathrm{colim}}\, map_{\mathscr{C}}(K_S, FX_{S,n}) \rightrightarrows \prod_{S\to T} \underset{\mathbb{N}}{\mathrm{colim}}\, map_{\mathscr{C}}(K_s, FX_{T,n}) \right) \end{aligned}$$

Here we used that finite limits and sequential colimits commute in $\mathscr{C}$ (see [AR94, 1.59]), and that the natural isomorphism in question is G-equivariant. Replacing the maps $X_{T,n} \to X_{T,n+1}$ and using that sequential colimits along cofibrations preserve fibrant objects (see [Sch13, 1.3.2]), we are left with showing that

$$\underset{\mathbb{N}}{\mathrm{colim}}(map_{\mathscr{C}}(K, Y_n)) \cong map_{\mathscr{C}}(K, \underset{\mathbb{N}}{\mathrm{colim}}\, Y_n)$$

for every finite simplicial set K and functor $Y\colon \mathbb{N} \to \mathscr{C}$. By the Yoneda lemma and local presentability of $\mathscr{C}$, this follows from the sequence of natural bijections for an object c of $\mathscr{C}$.

$$\begin{aligned}\mathscr{C}(c, map_{\mathscr{C}}(K, \operatorname*{colim}_{\mathbb{N}} Y_n)) \cong \mathscr{C}(K \otimes c, \operatorname*{colim}_{\mathbb{N}} Y_n) \cong sSet(K, Map_{\mathscr{C}}(c, \operatorname*{colim}_{\mathbb{N}} Y_n)) \cong \\ sSet(K, \operatorname*{colim}_{\mathbb{N}} Map_{\mathscr{C}}(c, Y_n)) \cong \operatorname*{colim}_{\mathbb{N}} sSet(K, Map_{\mathscr{C}}(c, Y_n)) \cong \\ \operatorname*{colim}_{\mathbb{N}} \mathscr{C}(c, map_{\mathscr{C}}(K, Y_n)) \cong \mathscr{C}(c, \operatorname*{colim}_{\mathbb{N}} map_{\mathscr{C}}(K, Y_n)).\end{aligned}$$

□

References

[AR94] Jiří Adámek and Jiří Rosický, *Locally presentable and accessible categories*, London Mathematical Society Lecture Note Series, vol. 189, Cambridge University Press, Cambridge, 1994. MR 1294136 (95j:18001)

[BK72] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin, 1972. MR 0365573 (51 #1825)

[Blu06] Andrew J. Blumberg, *Continuous functors as a model for the equivariant stable homotopy category*, Algebr. Geom. Topol. **6** (2006), 2257–2295. MR 2286026 (2008a:55006)

[DI04] Daniel Dugger and Daniel C. Isaksen, *Topological hypercovers and $\mathbb{A}^1$-realizations*, Math. Z. **246** (2004), no. 4, 667–689. MR 2045835 (2005d:55026)

[Dot13] E. Dotto, *Calculus of equivariant functors and $\mathbb{Z}/2$-analyticity of real k-theory*, arXiv: 1305.6217, 2013.

[DS07] Daniel Dugger and Brooke Shipley, *Enriched model categories and an application to additive endomorphism spectra*, Theory Appl. Categ. **18** (2007), No. 15, 400–439. MR 2342167 (2009d:18012)

[Dug01] Daniel Dugger, *Combinatorial model categories have presentations*, Adv. Math. **164** (2001), no. 1, 177–201. MR 1870516 (2002k:18022)

[Elm83] A. D. Elmendorf, *Systems of fixed point sets*, Trans. Amer. Math. Soc. **277** (1983), no. 1, 275–284. MR 690052 (84f:57029)

[GJ09] Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Modern Birkhäuser Classics, Birkhäuser Verlag, Basel, 2009, Reprint of the 1999 edition [MR1711612]. MR 2840650

[GM13] B. Guillou and J.P. May, *Enriched model categories and diagram categories*, arXiv: 1110.3567, 2013.

[Goo92] Thomas G. Goodwillie, *Calculus. II. Analytic functors*, K-Theory **5** (1991/92), no. 4, 295–332. MR 1162445 (93i:55015)

[Hir03] Philip S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99, American Mathematical Society, Providence, RI, 2003. MR 1944041 (2003j:18018)

[JS01] Stefan Jackowski and Jolanta Słomińska, *G-functors, G-posets and homotopy decompositions of G-spaces*, Fund. Math. **169** (2001), no. 3, 249–287. MR 1852128 (2002h:55017)

[Kel82] G. M. Kelly, *Structures defined by finite limits in the enriched context. I*, Cahiers Topologie Géom. Différentielle **23** (1982), no. 1, 3–42, Third Colloquium on Categories, Part VI (Amiens, 1980). MR 648793 (83h:18007)

[May75] J. Peter May, *Classifying spaces and fibrations*, Mem. Amer. Math. Soc. **1** (1975), no. 1, 155, xiii+98. MR 0370579 (51 #6806)

[MM02] M. A. Mandell and J. P. May, *Equivariant orthogonal spectra and S-modules*, Mem. Amer. Math. Soc. **159** (2002), no. 755, x+108. MR 1922205 (2003i:55012)

[Rie13] Emily Riehl, *Categorical homotopy theory*, Lecture notes, 2013.

[Sch13] Stefan Schwede, *Lectures on equivariant stable homotopy theory*, 2013.

[Shi03] Brooke Shipley, *A convenient model category for commutative ring spectra*, Tech. report, 2003.

[Ste10] Marc Stephan, *Elmendorf's theorem for cofibrantly generated model categories*, Master's thesis, ETH Zurich, 2010.

[Tho79] R. W. Thomason, *Homotopy colimits in the category of small categories*, Math. Proc. Cambridge Philos. Soc. **85** (1979), no. 1, 91–109. MR 510404 (80b:18015)

[TW91] J. Thvenaz and P. J. Webb, *Homotopy equivalence of posets with a group action*, J. Combin. Theory Ser. A **56** (1991), 173–181.

[VF99] Rafael Villarroel-Flores, *Equivariant homotopy type of categories and preordered sets*, PhD. Thesis, University of Minnesota, 1999.

[VF04] ———, *The action by natural transformations of a group on a diagram of spaces*, arXiv: 0411502v1, 2004.